

Datasheet for ABIN7317896 **ERBB4 Protein (Fc Tag)**



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	ERBB4
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This ERBB4 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Human/Rhesus HER4/ErbB4 Protein (Fc Tag)(Active)
Sequence:	Met 1-Arg649
Characteristics:	A DNA sequence encoding the human ERBB4 (NP_005226.1) (Met1-Arg649) was expressed, fused with the Fc region of human IgG1 at the C-terminus. Human and Rhesus ERBB4 sequences are identical.
Purity:	> 85 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	1. Measured by its ability to bind biotinylated human Fc-NGR1 (isoform Beta1) in a functional ELISA. 2. Measured by its ability to bind biotinylated human NRG1 (isoform Beta1) in a functional ELISA. 3. Measured by its ability to bind biotinylated human NRG1 (aa 2-246)-Fc in a functional ELISA.

Target Details

Target:	ERBB4
Alternative Name:	ErbB4 (ERBB4 Products)
Background:	<p>Background: ERBB4 is a single-pass type I membrane protein with multiple cysteine rich domains; a transmembrane domain; a tyrosine kinase domain; a phosphatidylinositol-3 kinase binding site and a PDZ domain binding motif. ERBB4 is expressed at highest levels in brain; heart; kidney; in addition to skeletal muscle; parathyroid; cerebellum; pituitary; spleen; testis and breast. And lower levels in thymus; lung; salivary gland; and pancreas. It specifically binds to and is activated by neuregulins; NRG-2; NRG-3; heparin-binding EGF-like growth factor; betacellulin and NTAK. ERBB4 also can be activated by other factors and induces a variety of cellular responses including mitogenesis and differentiation. ERBB4 regulates development of the heart; the central nervous system and the mammary gland; gene transcription; cell proliferation; differentiation; migration and apoptosis. It is required for normal cardiac muscle differentiation during embryonic development; and for postnatal cardiomyocyte proliferation. ERBB4 also play a role on the normal development of the embryonic central nervous system; especially for normal neural crest cell migration and normal axon guidance. It is required for mammary gland differentiation; induction of milk proteins and lactation.</p> <p>Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy</p> <p>Synonym: Receptor tyrosine-protein kinase erbB-4; Proto-oncogene-like protein c-ErbB-4; Tyrosine kinase-type cell surface receptor HER4; p180erbB4; ERBB4; HER4</p>
Molecular Weight:	96.6 kDa
NCBI Accession:	NP_005226
Pathways:	RTK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway

Application Details

Comment:	117 kDa
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4

Handling

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.