

Datasheet for ABIN7317909 ICAM2 Protein (His tag,Fc Tag)



OverviewQuantity:100 µgTarget:ICAM2Origin:HumanSource:HEK-293 CellsProtein Type:RecombinantBiological Activity:ActivePurification tag / Conjugate:This ICAM2 protein is labelled with His tag,Fc Tag.

Product Details

Purpose:	Recombinant Human ICAM-2/CD102 Protein (His & Fc Tag)(Active)
Sequence:	Met 1-Gln 223
Characteristics:	A DNA sequence encoding the extracellular domain (Met 1-Gln 223) of human ICAM2 (NP_000864.2) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Purity:	> 97 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by the ability of the immobilized protein to support the adhesion of PMA-stimulated HSB2 human peripheral blood acute lymphoblastic leukemia cells. When cells are added to ICAM2-coated plates (12.5 μ g/mL, 100 μ L/well), approximately 50 % - 60 % will adhere specifically.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7317909 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Target:	ICAM2
Alternative Name:	ICAM-2/CD102 (ICAM2 Products)
Background:	Background: Intercellular adhesion molecule 2 (ICAM-2, CD102), belongs to the ICAM family consisting of three members identified as ligands for integrin receptors. It is a type I transmembrane glycoprotein with two lg-like C2-type domains, and binds to the leukocyte integrins LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18). As a second ligand of leukocyte function-associated antigen-1, ICAM-2 functions as a costimulatory molecule for effector cells. ICAM-2 is mainly expressed on vascular endothelial and hematopoietic cells. Interactions of ICAM-2 and the integrin receptors mediate cell adhesion in a wide range of lymphocyte, monocyte, natural killer cell, and granulocytewith other cells, and play important roles in many adhesion-dependent immune and inflammation responses, such as T cell aggregation, NK-cell cytotoxicity and migration, lymphocyte recirculation, etc. Serum levels of ICAM-2 correlated significantly with the inflammatory and course sequences of trichinosis in mice and had a similar relation with blood eosinophilia. So, estimation of ICAM-2 serum levels may prove useful in diagnosis of trichinosis recent infections, and in monitoring the prognosis and response to treatment. Synonym: Intercellular Adhesion Molecule 2, ICAM-2, CD102, ICAM2
Molecular Weight:	50.3 kDa
NCBI Accession: Application Details	NP_000864
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN7317909 | 07/25/2024 | Copyright antibodies-online. All rights reserved.