

Datasheet for ABIN7317946
MMP8 Protein (His tag)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	MMP8
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This MMP8 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human MMP8/CLG1 Protein (His Tag)(Active)
Sequence:	Met 1-Gly 467
Characteristics:	A DNA sequence encoding the human MMP8 (NP_002415.1) (Met 1-Gly 467) was expressed, fused with a polyhistidine tag at the C-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to cleave the fluorogenic peptide substrate, Mca-PLGL-Dpa-AR-NH2 (AnaSpec, Catalog#27076). The specific activity is > 250 pmoles/min/µg.(Activation description: The proenzyme needs to be activated by APMA for an activated form)

Target Details

Target:	MMP8
---------	------

Target Details

Alternative Name:	MMP8/CLG1 (MMP8 Products)
Background:	<p>Background: Matrix metalloproteinases (MMPs) are a family of zinc-dependent endopeptidases that degrade components of the extracellular matrix (ECM) and play essential roles in various physiological processes such as morphogenesis, differentiation, angiogenesis and tissue remodeling, as well as pathological processes including inflammation, arthritis, cardiovascular diseases, pulmonary diseases and tumor invasion. Neutrophil collagenase, also known as Matrix metalloproteinase-8, MMP-8, and CLG1, is a member of the peptidase M10A family. MMP-8 may affect the metastatic behaviour of breast cancer cells through protection against lymph node metastasis, underlining the importance of anti-target identification in drug development. MMP-8 in the tumour may have a protective effect against lymph node metastasis. MMP-8 may affect the metastatic behaviour of breast cancer cells through protection against lymph node metastasis, underlining the importance of anti-target identification in drug development. MMP-8 participates in wound repair by contributing to the resolution of inflammation and open the possibility to develop new strategies for treating wound healing defects.</p> <p>Synonym: Neutrophil collagenase, Matrix metalloproteinase-8, MMP-8, PMNL collagenase, PMNL-CL,MMP8, CLG1</p>
Molecular Weight:	52.6 kDa
NCBI Accession:	NP_002415

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 50 mM Tris, 10 mM CaCl ₂ , 150 mM NaCl, pH 7.5
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.