

Datasheet for ABIN7317963 Layilin Protein (LAYN) (Fc Tag)



Overview	
Quantity:	100 µg
Target:	Layilin (LAYN)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Layilin protein is labelled with Fc Tag.
Product Details	
Purpose:	Recombinant Human Layilin/LAYN Protein (Fc Tag)
Sequence:	Met 1-Glu 220
Characteristics:	A DNA sequence encoding the extracellular domain (Met 1-Glu 220) of human Layilin
	(NP_849156.1) precursor was expressed with the C-terminal fused Fc region of human IgG1.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per μ g as determined by the LAL method.
Target Details	
Target:	Layilin (LAYN)
Alternative Name:	Layilin/LAYN (LAYN Products)
Background:	Background: Layilin, a recently characterized as a 55 kDa transmembrane protein with
	homology to C-type lectins, is present in numerous cell lines and tissue extracts. As one of the
	adaptor proteins, talin mediates the interactions between the actin filaments and the cell
Order at www.antibo	odies-online.com www.antikoerper-online.de www.anticorps-enligne.fr www.antibodies-online.cn

International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7317963 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

membrane by binding to integral membrane proteins and to the cytoskeleton. Layilin is a newly
identified membrane-binding site for talin in peripheral ruffles of spreading cells, a ten-amino
acid motif in the layilin cytoplasmic domain is sufficient for talin binding, and its adjacent LH2-
LH3 tandem arrays in the cytoplasmic domain provide docking sites for talin. Furthermore, talin
binds layilin, PIPK1gamma and integrins in similar although subtly different ways. Layilin binds
specifically to hyaluronan (HA) through its extracellular domain, a ubiquitous extracellular
matrix component in most animal tissues and body fluids, but not to other tested
glycosaminoglycans. The research suggests that layilin may mediate signals from extracellular
matrix to the cell cytoskeleton via interaction with different intracellular binding partners and
thereby be involved in the modulation of cortical structures in the cell. All the above actions
reveal an interesting parallel between layilin and the known HA receptor CD44. In addition,
merlin and radixin have been identified as different intracellular binding partners of layilin.
Accordingly, it has been suggested that layilin plays roles in a variety of cellular processes,
including cell shape, adhesion, motility, and homeostasis, as well as signal transduction. In
addition, layilin might play an important role in the process of invasion and lymphatic
metastasis of lung carcinoma.
Synonym: Layilin

Molecular Weight:	49.5 kDa
NCBI Accession:	NP_849156

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 100 mM Glycine, 10 mM NaCl, 50 mM Tris, pH 7.5
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN7317963 | 07/25/2024 | Copyright antibodies-online. All rights reserved.