

[Go to Product page](#)

Datasheet for ABIN7318040 CD55 Protein (Fc Tag)

Overview

Quantity:	100 µg
Target:	CD55
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD55 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Human CD55/DAF Protein (Fc Tag)(Active)
Sequence:	Met 1-Ser 353
Characteristics:	A DNA sequence encoding the human CD55 precursor (NP_000565.1) (Met 1-Ser 353) was expressed with the C-terminal fused Fc region of human IgG1.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized recombinant human CD97 at 10 µg/ml (100 µl/well) can bind recombinant human CD55 at a linear range of 0.46-30 µg/ml.

Target Details

Target:	CD55
---------	------

Target Details

Alternative Name: CD55/DAF ([CD55 Products](#))

Background: CD55, also well known as decay-accelerating factor (DAF), is a member of the RCA (regulators of complement activation) family characterized by four to 30 SCRs (short consensus repeats) in their plasma-exposed regions. It is a major regulator of the alternative and classical pathways of complement activation and is expressed on all serum-exposed cells. CD55 is physiologically acting as an inhibitor of the complement system, but is also broadly expressed in malignant tumours. DAF seems to exert different functions beyond its immunological role such as promotion of tumorigenesis, decrease of complement mediated tumor cell lysis, autocrine loops for cell rescue and evasion of apoptosis, neoangiogenesis, invasiveness, cell motility. It is commonly hijacked by invading pathogens, including many enteroviruses and uropathogenic Escherichia coli, to promote cellular attachment prior to infection. This 70-75 kDa glycoprotein CD55 containing four SCR modules is involved in the regulation of the complement cascade. It inhibits complement activation by suppressing the function of C3/C5 convertases, thereby limiting local generation or deposition of C3a/C5a and membrane attack complex (MAC or C5b-9) production. DAF has been identified as a ligand for an activation-associated, seven-transmembrane lymphocyte receptor, CD97, which is a receptor mediating attachment and infection of several viruses and bacteria. In addition, it has been shown that DAF regulates the interplay between complement and T cell immunity in vivo, and thus may be implicated in immune and tumor biology.

Synonym: Complement Decay-Accelerating factor, CD55, CR,CROM,DAF,TC

Molecular Weight: 61.7 kDa

NCBI Accession: [NP_000565](#)

Pathways: [Complement System](#), [Regulation of Leukocyte Mediated Immunity](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C, -20 °C, -80 °C

Handling

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.