



[Go to Product page](#)

Datasheet for ABIN7318104
EPH Receptor A2 Protein (EPHA2) (His tag)

Overview

Quantity:	50 µg
Target:	EPH Receptor A2 (EPHA2)
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This EPH Receptor A2 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human EphA2 Protein (His Tag)(Active)
Sequence:	Ala24-Asn534
Characteristics:	Recombinant Human Ephrin A Receptor 2 is produced by our Mammalian expression system and the target gene encoding Ala24-Asn534 is expressed with a 6His tag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Immobilized Human EphA2-His at 4µg/ml(100 µl/well) can bind Human EFNA1-Fc(Cat: PKSH032389). The ED50 of Human EphA2-His is 0.03ug/ml .

Target Details

Target:	EPH Receptor A2 (EPHA2)
---------	-------------------------

Target Details

Alternative Name: EphA2 ([EPHA2 Products](#))

Background: Eph receptor A2 (Ephrin type-A receptor 2 or EphA2) is a member of the ephrin receptor subfamily of the protein-tyrosine kinase family. The Eph receptors' corresponding family of ligands are the ephrins anchored to cell surfaces. The ephrins and Eph receptors are implicated as positional labels that may guide the development of neural topographic maps. They have also been found implicated in embryonic patterning, neuronal targeting, vascular development and adult neovascularization. The large family of ligands and receptors may make a major contribution to the accurate spatial patterning of connections and cell position in the nervous system. Furthermore, elevated expression of Eph receptors and ephrin ligands is associated with tumors and associated tumor vasculature, suggesting the Eph receptors and ephrin ligands also play critical roles in tumor angiogenesis and tumor growth. Unlike most Eph kinases, which are primarily expressed during development, EphA2 is primarily found in adult human epithelial cells. The cellular functions of EphA2 may be regulating cell growth, survival, migration, and angiogenesis. Unlike other receptor tyrosine kinases, ligand binding is not necessary for EphA2. Rather, the ligand appears to regulate EphA2 subcellular localization and its interactions with downstream adapter and signaling proteins. Eph receptor A2(EphA2) has been demonstrated to critically regulate tumor cell growth, migration and invasiveness. Eph receptor A2(EphA2) is frequently overexpressed and functionally altered in aggressive tumor cells, and that these changes promote metastatic character.

Synonym: Ephrin type-A receptor 2, Epithelial cell kinase, Tyrosine-protein kinase receptor ECK, EPHA2,ARCC2,CTPA,CTPP1,CTRCT6,ECK

Molecular Weight: 57.0 kDa

UniProt: [P29317](#)

Pathways: [RTK Signaling](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Handling

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.