

Datasheet for ABIN7318578

HDGF Protein (His tag)



| _ | | | | | |
|---|-----|---|----|-------------|-----|
| | 1// | r | Vİ | \triangle | ۸/ |
| | V | | VI | | / V |

| Quantity: | 50 µg |
|-------------------------------|---|
| Target: | HDGF |
| Origin: | Human |
| Source: | Escherichia coli (E. coli) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This HDGF protein is labelled with His tag. |
| Product Details | |
| Purpose: | Recombinant Human HDGF Protein (His Tag) |
| Sequence: | Met 1-Tyr100 |
| Characteristics: | Recombinant Human Hepatoma-Derived Growth Factor is produced by our E.coli expression system and the target gene encoding Met1-Tyr100 is expressed with a 6His tag at the C-terminus. |
| Purity: | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin Level: | < 1.0 EU per µg as determined by the LAL method. |
| Target Details | |
| Target: | HDGF |
| Alternative Name: | HDGF (HDGF Products) |
| Background: | Background: Hepatoma-Derived Growth Factor is a original member of the HDGF family. HDGF is a cytoplasmic protein and contains one PWWP domain. HDGF expression levels are high in |

the nucleus and cytoplasm of smooth muscle and endothelial cells. HDGF has proliferative, angiogenic and neurotrophic activity. HDGF was initially characterized as a secreted mitogen from the Huh-7 human hepatoma cell line. As a heparin-binding protein, which is highly expressed in tumor cells where it stimulates proliferation. HDGF has mitogenic activity for fibroblasts and acts as a transcriptional repressor. It has been shown that HDGF is linked with tumorigenesis and the growth of cancer.

Synonym: Hepatoma-Derived Growth Factor, HDGF, High Mobility Group Protein 1-Kike 2, HMG-1L2, HDGF, HMG1L2

Molecular Weight: 12.6 kDa

UniProt: P51858

Pathways: ER-Nucleus Signaling

Application Details

Restrictions: For Research Use only

Handling

| Format: | Lyophilized | | |
|------------------|---|--|--|
| Reconstitution: | Please refer to the printed manual for detailed information. | | |
| Buffer: | Lyophilized from a 0.2 µm filtered solution of 20 mM Tris, 1 mM DTT, 1 mM EDTA, pH 7.5. | | |
| Storage: | 4 °C,-20 °C,-80 °C | | |
| Storage Comment: | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. | | |
| | Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted | | |
| | samples are stable at < -20°C for 3 months. | | |