# antibodies -online.com





# IL-18 Protein (His tag)



| O       | :   |
|---------|---|
| Over    | $\backslash \backslash \backslash \triangle \backslash \backslash \backslash$ |
| $\circ$ | V I C V V   |

| 0.0                           |  |
|-------------------------------|--|
| Quantity:                     | 50 μg  |
| Target:                       | IL-18 (IL18)   |
| Origin:                       | Human  |
| Source:                       | Human Cells  |
| Protein Type:                 | Recombinant  |
| Purification tag / Conjugate: | This IL-18 protein is labelled with His tag.   |
| Product Details               |  |
| Purpose:                      | Recombinant Human Interleukin-18/IL-18 Protein (His Tag)   |
| Sequence:                     | Tyr37-Asp193   |
| Characteristics:              | Recombinant Human Interleukin-18 is produced by our Mammalian expression system and the target gene encoding Tyr37-Asp193 is expressed with a 6His tag at the C-terminus.  |
| Purity:                       | > 95 % as determined by reducing SDS-PAGE.   |
| Endotoxin Level:              | < 1.0 EU per µg as determined by the LAL method.   |
| Target Details                |  |
| Target:                       | IL-18 (IL18)   |
| Alternative Name:             | Interleukin-18/IL-18 (IL18 Products)   |
| Background:                   | Background: Interleukin-18 is a secreted protein and it belongs to the IL-1 family. IL-18 is a proinflammatory cytokine and produced by macrophages and other cells. This cytokine can induce the IFN-gamma production of T cells. The combination of this cytokine and IL12 has |

#### **Target Details**

been shown to inhibit IL-4 dependent IgE and IgG1 production, and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity. After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon-γ (IFN-γ) or type II interferon that plays an important role in activating the macrophages or other cells. Synonym: Interleukin-18, Iboctadekin, Interferon gamma-inducing factor, IFN-gamma-inducing factor, Interleukin-1 gamma, IL-1 gamma, GIF, IL-18, IL-1g, IL1F4, MGC12320

Molecular Weight: 23.3 kDa

UniProt: Q14116

Pathways: Cellular Response to Molecule of Bacterial Origin, Activated T Cell Proliferation, Cancer Immune

Checkpoints, Inflammasome

## **Application Details**

Restrictions: For Research Use only

## Handling

| Format:          | Lyophilized  |
|------------------|--|
| Reconstitution:  | Please refer to the printed manual for detailed information.   |
| Buffer:          | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.  |
| Storage:         | 4 °C,-20 °C,-80 °C   |
| Storage Comment: | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.  Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted |

samples are stable at < -20°C for 3 months.