antibodies .- online.com







IDH1 Protein (Arg132His-Mutant) (His tag)



Overview

50 µg
IDH1
Arg132His-Mutant
Human
Escherichia coli (E. coli)
Recombinant
This IDH1 protein is labelled with His tag.
Recombinant Human IDH1 Protein (R132H, C-8His)
Met1-Leu414(Arg132His)
Recombinant Human Isocitrate Dehydrogenase [NADP] Cytoplasmic is produced by our E.coli
expression system and the target gene encoding Met1-Leu414(Arg132His) is expressed with a
expression system and the target gene encoding Met1-Leu414(Arg132His) is expressed with a
expression system and the target gene encoding Met1-Leu414(Arg132His) is expressed with a 8His tag at the C-terminus.
expression system and the target gene encoding Met1-Leu414(Arg132His) is expressed with a 8His tag at the C-terminus. > 95 % as determined by reducing SDS-PAGE.
expression system and the target gene encoding Met1-Leu414(Arg132His) is expressed with a 8His tag at the C-terminus. > 95 % as determined by reducing SDS-PAGE.

Target Details

Dackground.	Bac	karo	und:
-------------	-----	------	------

Background: Isocitrate Dehydrogenase [NADP] Cytoplasmic (IDH1) belongs to the isocitrate and isopropylmalate dehydrogenases family. IDH1 exists as a homodimer, binding one magnesium or manganese ion per subunit. Mutations of IDH1 have been shown to cause metaphyseal chondromatosis with aciduria and are involved in the development of glioma. IDH plays a role in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the α-hydroxylation of phytanic acid.

Synonym: Isocitrate Dehydrogenase [NADP] Cytoplasmic, IDH, Cytosolic NADP-Isocitrate Dehydrogenase, IDP, NADP(+)-Specific ICDH, Oxalosuccinate Decarboxylase, IDH1, PICD

Molecular Weight:

48.1 kDa

UniProt:

075874

Pathways:

Warburg Effect

Application Details

Restrictions:

For Research Use only

Handling

Format:	Frozen, Liquid
Buffer:	Supplied as a 0.2 µm filtered solution of 50 mM Tris, 200 mM NaCl, 10 % glycerol, pH 8.0.
Storage:	-20 °C
Storage Comment:	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.