

Datasheet for ABIN7319094

Osteoprotegerin Protein (Fc Tag)



Go to Product page

()	11/	\sim	r۱	۲i	\cap	۱۸/	ı
\cup	V	L	۲V	1	ヒ	٧V	

Quantity:	50 μg
Target:	Osteoprotegerin (TNFRSF11B)
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Osteoprotegerin protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Human Osteoprotegerin/TNFRSF11B Protein (Fc Tag)(Active)
Sequence:	Glu22-Leu201
Characteristics:	Recombinant Human Osteoprotegerin is produced by our Mammalian expression system and the target gene encoding Glu22-Leu201 is expressed with a Fc tag at the C-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Immobilized Human OPG-Fc at $2\mu g/ml(100\ \mu l/well)$ can bind Mouse RANKL-His(Cat: PKSM041165). The ED50 of Mouse OPG-Fc is $2.44ug/ml$.

Target Details

	Osteoprotegerin (TNFRSF11B)	Target:
--	-----------------------------	---------

Target Details

Storage Comment:

rarget Details	
Alternative Name:	Osteoprotegerin/TNFRSF11B (TNFRSF11B Products)
Target Type:	Chemical
Background:	Background: TNFRSF11B is a secreted protein, containing 2 death domains and 4 TNFR-Cys
	repeats. TNFRSF11B is a decoy receptor for the receptor activator of nuclear factor kappa B
	ligand (RANKL). By binding RANKL, TNFRSF11B inhibits nuclear kappa B (NF-кВ) which is a
	central and rapid acting transcription factor for immune-related genes, and a key regulator of
	inflammation, innate immunity, and cell survival and differentiation. TNFRSF11B levels are
	influenced by voltage-dependent calcium channelsCav1.2. TNFRSF11B can reduce the
	production of osteoclasts by inhibiting the differentiation of osteoclast precursors into
	osteoclasts and also regulates the resorption of osteoclasts in vitroand in vivo. TNFRSF11B
	binding to RANKL on osteoblast/stromal cells, blocks the RANKL-RANK ligand interaction
	between osteoblast/stromal cells and osteoclast precursors. This has the effect of inhibiting
	the differentiation of the osteoclast precursor into a mature osteoclast.
	Synonym: Tumor necrosis factor receptor superfamily member 11B, Osteoclastogenesis
	inhibitory factor, Osteoprotegerin, TNFRSF11B, OCIF, OPG,PDB5,TR1
Molecular Weight:	47.2 kDa
UniProt:	000300
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C

samples are stable at < -20°C for 3 months.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted