## .-online.com antibodies

# Datasheet for ABIN7319188 **ZBTB9 Protein (His tag)**



#### Overview

Quantity:	50 µg
Target:	ZBTB9
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ZBTB9 protein is labelled with His tag.

#### Product Details

Purpose:	Recombinant Human ZBTB9 Protein (His Tag)
Sequence:	Met 1-Lys473
Characteristics:	Recombinant Human Zinc Finger and BTB domain-containing protein 9 is produced by our E.coli expression system and the target gene encoding Met1-Lys473 is expressed with a 6His tag at the C-terminus.
Purity:	> 80 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per $\mu$ g as determined by the LAL method.

### Target Details

Target:	ZBTB9
Alternative Name:	ZBTB9 (ZBTB9 Products)
Background:	Background: Zinc finger and BTB domain-containing protein 9 is a protein-coding gene. ZBTB9
	is a 473 amino acids protein that Contains 1 BTB (POZ) domain and 2 C2H2-type zinc fingers.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7319188 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	Diseases associated with ZBTB9 include systemic lupus erythematosus, and lupus erythematosus. Zinc finger and BTB domain-containing protein 9 may be involved in transcriptional regulation. Synonym: Zinc finger and BTB domain-containing protein 9,ZBTB9
Molecular Weight:	51.7 kDa
UniProt:	Q96C00
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted