.-online.com antibodies

Datasheet for ABIN7319207 SNCA Protein (His tag)



Overview	
Quantity:	50 µg
Target:	SNCA
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SNCA protein is labelled with His tag.
Product Details	
Purpose:	Recombinant Human Alpha-Synuclein/SNCA Protein (His Tag)
Sequence:	Met 1-Ala140
Characteristics:	Recombinant Human alpha-Synuclein is produced by our E.coli expression system and the target gene encoding Met1-Ala140 is expressed with a 6His tag at the N-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Characteristics: Purity:	Recombinant Human alpha-Synuclein is produced by our E.coli expression system and the target gene encoding Met1-Ala140 is expressed with a 6His tag at the N-terminus.

Target Details

Target:	SNCA
Alternative Name:	Alpha-Synuclein/SNCA (SNCA Products)
Background:	Background: Alpha-Synuclein (SNCA) is a member of the Synuclein family. SNCA is expressed principally in brain but also expressed in low concentrations in all tissues except liver. SNCA
	interacts with UCHL1, Phospholipase D and histones. SNCA can include beta- and gamma-

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7319207 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	synuclein. In addition, SNCA is an important regulatory component of vesicular transport in
	neuronal cells. It has been suggested that SNCA is related to the pathogenesis of Parkinson's
	Disease and neurodegenerative disorders. Defects in SNCA will lead to Dementia Lewy Body
	(DLB).
	Synonym: Alpha-Synuclein, Non-A Beta Component of AD Amyloid, Non-A4 Component of
	Amyloid Precursor, NACP, SNCA, NACP, PARK1
Molecular Weight:	16.7 kDa
UniProt:	P37840
Pathways:	Synaptic Membrane, Regulation of G-Protein Coupled Receptor Protein Signaling, Positive
	Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process, Platelet
	derived growth Factor Receptor Signaling, Negative Regulation of Transporter Activity,
	Regulation of long-term Neuronal Synaptic Plasticity
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.