

Datasheet for ABIN7319329

R-Spondin 3 Protein (RSPO3) (His tag,Fc Tag)



Go to Product page

| () | ve | r\/i | | ۱۸/ |
|--------|-----|-------|---------------|-----|
| \cup | V C | 1 / 1 | $\overline{}$ | ٧V |

Background:

| Quantity: | 50 μg |
|-------------------------------|--|
| Target: | R-Spondin 3 (RSP03) |
| Origin: | Human |
| Source: | Human Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This R-Spondin 3 protein is labelled with His tag,Fc Tag. |
| Product Details | |
| Purpose: | Recombinant Human RSP03 Protein (Fc & His Tag) |
| Sequence: | Gln22-His272 |
| Characteristics: | Recombinant Human R-spondin-3 is produced by our Mammalian expression system and the target gene encoding Gln22-His272 is expressed with a Fc, 6His tag at the C-terminus. |
| Purity: | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin Level: | < 1.0 EU per µg as determined by the LAL method. |
| Target Details | |
| Target: | R-Spondin 3 (RSP03) |
| Alternative Name: | RSP03 (RSP03 Products) |

Background: R-spondin-3 (RSPO3), also known as Protein with TSP type-1 repeat, Roof plate-specific spondin-3, Thrombospondin type-1 domain-containing protein 2, PWTSR, THSD2 and

CRISTIN1, is a member of the thrombospondin type 1 repeat supergene family. RSP03 is a

Target Details

secreted protein and widely expressed in many tissues. RSPO3 contains two Furin-like repeats which have been found in a variety of eukaryotic proteins involved in the mechanism of signal transduction by receptor tyrosine kinases, and one TSP type-1 domain, RSPO3 founctions as a activator of the beta-catenin signaling cascade, leading to TCF-dependent gene activation. Otherwise, RSPO3 may negatively regulate the TGF-beta pathway.

Synonym: R-spondin-3,RSPO3,Protein with TSP type-1 repeat, Roof plate-specific spondin-3, Thrombospondin type-1 domain-containing protein 2, PWTSR, THSD2, CRISTIN1

Molecular Weight:

56.3 kDa

Application Details

Restrictions:

For Research Use only

Handling

| Format: | Lyophilized | |
|------------------|--|--|
| Reconstitution: | Please refer to the printed manual for detailed information. | |
| Buffer: | Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.2. | |
| Storage: | 4 °C,-20 °C,-80 °C | |
| Storage Comment: | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted | |
| | samples are stable at < -20°C for 3 months. | |