# antibodies -online.com





## CD226 Protein (CD226) (Fc Tag)



GO to Product page

( )	11/0	K\ /	iew
	$\cup$	ועוי	$I \cap VV$

**Product Details** 

Quantity:	50 μg
Target:	CD226
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD226 protein is labelled with Fc Tag.

1 Todas Betano		
Purpose:	Recombinant Human CD226/DNAM-1 Protein (Fc Tag)(Active)	
Sequence:	Glu19-Asn247	
Characteristics:	Recombinant Human CD226 Antigen is produced by our Mammalian expression system and the target gene encoding Glu19-Asn247 is expressed with a Fc tag at the C-terminus.	
Purity:	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.	
Biological Activity Comment: Immobilized Human DNAM-1-Fc at 2µg/ml(100 µl/well) can bind Human Nectin-2-6l PKSH032913). The ED50 of Human DNAM-1-Fc is 0.79 ug/mL		

## Target Details

|--|

### **Target Details**

rarget Details		
Alternative Name:	CD226/DNAM-1 (CD226 Products)	
Background:	Background: Human DNAX accessory molecule 1 (DNAM-1/CD226) is a 65 kDa type I	
	transmembrane glycoprotein in the immunoglobulin superfamily. Mature human DNAM-1	
	contains an extracellular domain (ECD) with two Ig-like C2-set domains and a cytoplasmic	
	region that contains motifs for binding PDZ domains and band 4.1 family proteins. DNAM-1 is	
	expressed on multiple lymphoid and myeloid cells and interacts with CD155 and CD112.	
	Ligation of DNAM-1 promotes the activation of NK cells, CD8+ T cells, and mast cells, dendritic	
	cell maturation, megakaryocyte and activated platelet adhesion to vascular endothelial cells,	
	and monocyte extravasation, it inhibits the forrmation of osteoclasts. Plateletendothelium,	
	interactions mediated by DNAM-1, enable the metastasis of tumor cells to the lung.	
	Synonym: CD226 antigen, DNAX accessory molecule 1, DNAM-1, CD226, DNAM1	
Molecular Weight:	52.8 kDa	
UniProt:	Q15762	
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process,	
	Cancer Immune Checkpoints	
Application Datails		
Application Details		
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	