

## Datasheet for ABIN7319702

# **AMIGO2 Protein (Fc Tag)**



#### Overview

| Quantity:                     | 50 μg  |
|-------------------------------|--|
| Target:                       | AMIGO2                                       |
| Origin:                       | Human  |
| Source:                       | Human Cells                                  |
| Protein Type:                 | Recombinant                                  |
| Purification tag / Conjugate: | This AMIGO2 protein is labelled with Fc Tag. |
|                               |  |

### **Product Details**

| Purpose:         | Recombinant Human AMIGO2 Protein (Fc Tag)   |
|------------------|---|
| Sequence:        | Gly39-His393  |
| Characteristics: | Recombinant Human AMIGO2 is produced by our Mammalian expression system and the target gene encoding Gly39-His393 is expressed with a Fc tag at the C-terminus. |
| Purity:          | > 95 % as determined by reducing SDS-PAGE.  |
| Endotoxin Level: | < 1.0 EU per µg as determined by the LAL method.  |

## Target Details

| Target:           | AMIGO2   |
|-------------------|--|
| Alternative Name: | AMIGO2 (AMIGO2 Products)   |
| Background:       | Background: Amphoterin-Induced Protein 2 (AMIGO2) is a single-pass type I membrane protein which belongs to the AMIGO family of immunoglobulin superfamily. Mature AMIGO2 contains |
|                   | an Ig-like C2-type (immunoglobulin-like) domain, 6 LRR (leucine-rich) repeats, a LRRCT domain,   |

as well as a LRRNT domain. AMIGO2 is mainly expressed in in breast, ovary, cervix, and uterus, although lower in lung, colon, and rectum. AMIGO2 required for depolarization-dependent survival of cultured cerebellar granule neurons. AMIGO2 may mediate homophilic as well as heterophilic cell-cell interaction with AMIGO1 or AMIGO3. AMIGO2 may contribute to signal transduction through its intracellular domain, and may be required for tumorigenesis of a subset of gastric adenocarcinomas.

Synonym: Amphoterin-Induced Protein 2, AMIGO-2, Alivin-1, Differentially Expressed in Gastric Adenocarcinomas, DEGA, ALI1

Molecular Weight:

67.5 kDa

UniProt:

086SJ2

### **Application Details**

Restrictions:

For Research Use only

#### Handling

| Format:          | Lyophilized   |
|------------------|---|
| Reconstitution:  | Please refer to the printed manual for detailed information.                                  |
| Buffer:          | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.                                   |
| Storage:         | 4 °C,-20 °C,-80 °C  |
| Storage Comment: | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.   |
|                  | Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted |
|                  | samples are stable at < -20°C for 3 months.   |