antibodies - online.com





MUC16 Protein (Biotin)



Image



\sim			
	IV/E	۱//۱۲	$I \cap V$

Quantity:	100 μg
Target:	MUC16 (CA125)
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MUC16 protein is labelled with Biotin.
Product Details	
Purpose:	Recombinant Human CA125/MUC16 (C-Fc-Avi) Biotinylated
Sequence:	Gly12660-Met12923
Characteristics:	Biotinylated Recombinant Human Mucin-16 is produced by our Mammalian expression system and the target gene encoding Gly12660-Met12923 is expressed with a Fc, Avi tag at the C-terminus.
Purity:	>95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Target Details	
Target:	MUC16 (CA125)
Alternative Name:	CA125 (CA125 Products)
Background:	Background: MUC16, also known as the CA125 antigen, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial

ovarian cancer therapy. MUC16 is over-expressed by tumor cells including ovarian and mesothelial cancers. The transmembrane form can adhere to mesothelin in the peritoneum, facilitating metastasis of ovarian cancer to the peritoneal cavity. MUC16 also binds galectin-1 on immune cells and enhances its expression on tumor cells. MUC16-expressing tumors adhere to NK cells, down-regulate CD16 and suppress NK response, which may promote immune evasion. MUC16 is also cyclically expressed in the endometrium and may contribute to immune privilege during pregnancy. In the eye, MUC16 and other mucins protect the cornea and keep it hydrated. It is altered on the conjunctival epithelium of patients with non-Sjogren dry eye syndrome.

Synonym: CA125 ovarian cancer antigen, CA125, CA-125, CA125MUC-16, FLJ14303, MUC16,mucin 16, cell surface associated, mucin-16

Molecular Weight:

57.3 kDa

UniProt:

Q8WXI7


Application Details

Comment: 80-120 kDa

Restrictions: For Research Use only

Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCl, 100 mM Glycine, pH 7.5.	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	



Western Blotting

Image 1.