# antibodies .- online.com





# SIGLEC10 Protein (mFc Tag)

2 Images



#### Overview

Quantity:	50 μg
Target:	SIGLEC10
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This SIGLEC10 protein is labelled with mFc Tag.

#### **Product Details**

Purpose:	Recombinant Human Siglec-10 (C-mFc)
Sequence:	Met17-Thr546
Characteristics:	Recombinant Human Sialic Acid-binding Ig-like Lectin 10 is produced by our Mammalian expression system and the target gene encoding Met17-Thr546 is expressed with a mFc tag at the C-terminus.
Purity:	>90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Immobilized Anti-Human Siglec10 mAb(Cat#NC035) at 2µg/ml (100 µl/well) can bind Human Siglec-10-mFc(Cat#C01G). The ED50 of Human Siglec-10-mFc(Cat#C01G) is 18.16 ng/ml.

#### **Target Details**


Target:	SIGLEC10

## Target Details

Alternative Name:	Siglec-10 (SIGLEC10 Products)
Background:	Background: Siglecs (sialic acid binding Ig-like lectins) are I-type lectins that belong to the
	immunoglobulin superfamily. They are characterized by an N-terminal Ig-like V-type domain
	which mediates sialic acid binding, followed by a varying number of Ig-like C2-type domains.
	Siglecs 5-11 constitute the CD33/Siglec-3 related group, and are differentially expressed in the
	hematopoietic system. Siglec-G is the apparent ortholog of human Siglec-10. We describe here
	a novel member of the siglec protein family that shares a similar structure including five Ig-like
	domains, a transmembrane domain, and a cytoplasmic tail containing two ITIM-signaling
	motifs. Siglec-10 was identified through database mining of an asthmatic eosinophil EST
	library. Siglec-10 binds sialated proteins and lipids in alpha 2,3 or alpha 2,6 linkage and shows a
	preference for GT1b gangliosides. This binding can be modulated by cis interactions of Siglec-
	10 with sialated molecules expressed on the same cell. When tyrosine phosphorylated, the
	cytoplasmic ITIMs interact with phosphatases SHP-1 and SHP-2 to propagate inhibitory
	signals. The Siglec-10-VAP-1 interaction seems to mediate lymphocyte adhesion to
	endothelium and has the potential to modify the inflammatory microenvironment via the
	enzymatic end products.
	Synonym: SIGLEC10, MGC126774, PRO940, Siglec10, SLG2, sialic acid-binding Ig-like lectin 10,
	Siglec-10,siglec-like gene 2, Siglec-like protein 2, SLG2sialic acid binding Ig-like lectin 10 Ig-like lectin 7
Molecular Weight:	84.6 kDa
UniProt:	Q96LC7
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted


samples are stable at < -20°C for 3 months.

#### **Images**



### **Western Blotting**

Image 1.



#### **ELISA**

Image 2.