

Datasheet for ABIN7319836

CD3 epsilon Protein (CD3E) (Biotin)



Overview

| Overview | |
|-------------------------------|--|
| Quantity: | 100 μg |
| Target: | CD3 epsilon (CD3E) |
| Origin: | Human |
| Source: | Human Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CD3 epsilon protein is labelled with Biotin. |
| Product Details | |
| Durnoco | Decembinant Human CD2 anailan/CD2E (C Ea Avi) Diatinylated |

| Purpose: | Recombinant Human CD3 epsilon/CD3E (C-Fc-Avi) Biotinylated |
|------------------|--|
| Sequence: | Asp23-Asp126 |
| Characteristics: | Biotinylated Recombinant Human T-cell Surface Glycoprotein CD3 Epsilon Chain is produced by our Mammalian expression system and the target gene encoding Asp23-Asp126 is expressed with a Fc, Avi tag at the C-terminus. |
| Purity: | >95 % as determined by reducing SDS-PAGE. |
| Endotoxin Level: | < 1.0 EU per µg as determined by the LAL method. |

Target Details

| Target: | CD3 epsilon (CD3E) |
|-------------------|---|
| Alternative Name: | CD3E (CD3E Products) |
| Background: | Background: T-Cell Surface Glycoprotein CD3 ε Chain (CD3ε) is a single-pass type I membrane |
| | protein. CD3ɛ contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3ɛ is a |

polypeptide encoded by the CD3E gene on chromosome 11 in humans. The T cell receptor-CD3 complex (TCR/CD3 complex) is involved in T-cell development and several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs). TCR/CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

Synonym: T-Cell Surface Glycoprotein CD3 Epsilon Chain, T-Cell Surface Antigen T3/Leu-4 Epsilon Chain, CD3e, CD3E, T3E

Molecular Weight: 40.5 kDa

UniProt: P07766

Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway

Application Details

Comment: 45-60 kDa

Restrictions: For Research Use only

Handling

Format:LyophilizedReconstitution:Please refer to the printed manual for detailed information.Buffer:Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.Storage:4 °C,-20 °C,-80 °CStorage Comment:Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.