.-online.com antibodies

Datasheet for ABIN7319888 BTN2A2 Protein (Biotin)

Image



Overview

Quantity:	100 µg
Target:	BTN2A2
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This BTN2A2 protein is labelled with Biotin.

Product Details

Purpose:	Recombinant Human Butyrophilin Subfamily 2 Member A2/BTN2A2 (C-Fc-Avi) Biotinylated
Sequence:	Gln33-Val237
Characteristics:	Biotinylated Recombinant Human Butyrophilin Subfamily 2 Member A2 is produced by our Mammalian expression system and the target gene encoding Gln33-Val237 is expressed with a Fc, Avi tag at the C-terminus.
Purity:	>95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	BTN2A2
Alternative Name:	BTN2A2 (BTN2A2 Products)
Background:	Background: Butyrophilin 2A2 (BTN2A2) is a widely expressed type I transmembrane
	glycoprotein that functions as a negative regulator of immune responses. Mature human

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7319888 | 09/09/2023 | Copyright antibodies-online. All rights reserved.


	Butyrophilin 2A2 consisits of a 233 amino acid (aa) extracellular domain with two
	immunoglobulin-like domains, a 21 aa transmembrane segment, and a 237 aa cytoplasmic
	domain. Alternative splicing generates additional isoforms of human Butyrophilin 2A2 that lack
	the first, second, or both Iglike domains as well as isoforms with substitutions and deletions in
	the cytoplasmic region. Within the immune system, Butyrophilin 2A2 is expressed on thymic
	epithelial cells, naive B cells, splenic NK cells, dendritic cells, and peritoneal macrophages and is
	up-regulated with cell activation. Butyrophilin 2A2 inhibits T cell proliferation and activation and
	enhances the development of FoxP3+ regulatory T cells. Its up-regulation in the hippocampus
	is associated with schizophrenia.
	Synonym: Butyrophilin subfamily 2 member A2, BTN2A2
Molecular Weight:	51.5 kDa
UniProt:	Q8WVV5
Pathways:	Activated T Cell Proliferation

Application Details

Comment:	55-80 kDa
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7319888 | 09/09/2023 | Copyright antibodies-online. All rights reserved.