# antibodies -online.com





# LAIR1 Protein (AA 22-133) (His tag)





Go to Product page

$\sim$							
	1//	$\Box$	$r \setminus$	/ [	$\bigcirc$	1	٨,

Quantity:	100 μg
Target:	LAIR1
Protein Characteristics:	AA 22-133
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This LAIR1 protein is labelled with His tag.

### **Product Details**

Purpose:	Recombinant Mouse LAIR1 Protein (aa 22-133, His Tag)	
Sequence:	Gln 22-Ser 133	
Characteristics:	A DNA sequence encoding the mouse LAIR1 (Q8BG84-1) extracellular domain (Gln 22-Ser 133) was expressed, with a C-terminal polyhistidine tag.	
Purity:	> 90 % as determined by SDS-PAGE	
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.	

## **Target Details**

Target:	LAIR1
Alternative Name:	LAIR1 (LAIR1 Products)
Background:	Background: Leukocyte associated Ig-like receptor-1 (LAIR1) is a surface molecule expressed

on human mononuclear leukocytes that functions as an inhibitory receptor on human NK cells. In addition to NK cells, LAIR1 is expressed on T cells, B cells, macrophages, and dendritic cells. It is predicted to mediate inhibitory functions based on the presence of immunoreceptor tyrosine-based inhibitory motifs (ITIMs) in its cytoplasmic domain. Cross-linking of LAIR1 on human T cell clones results in inhibition of cytotoxicity only in T cell clones that lack CD28 and are able to spontaneously lyse certain targets in vitro. Moreover, the cytolytic activity of freshly isolated T cells, which is thought to be mainly due to "effector" T cells, can be inhibited by anti-LAIR1 mAb. Thus, LAIR1 functions as an inhibitory receptor not only on NK cells, but also on human T cells. This indicates that LAIR1 provides a mechanism of regulation of effector T cells and may play a role in the inhibition of unwanted bystander responses mediated by Ag-specific T cells.

Synonym: Leukocyte-associated immunoglobulin-like receptor 1; LAIR-1; mLAIR-1; CD305; Lair1

Molecular Weight:

14 kDa


#### **Application Details**

Restrictions:

For Research Use only

#### Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	



# **Western Blotting**

Image 1.