

Datasheet for ABIN7320078

VEGFR2/CD309 Protein (Fc Tag)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	VEGFR2/CD309 (VEGFR2)
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This VEGFR2/CD309 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Mouse VEGFR2/Flk-1/KDR Protein (Fc Tag)(Active)
Sequence:	Met1-Glu762
Characteristics:	A DNA sequence encoding the mouse KDR (P35918) (Met1-Glu762) was expressed, fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Immobilized mouse VEGF164 at 10 µg/ml (100 µl/well) can bind mouse KDR-Fc, The EC50 of mouse KDR-Fc is 0.11-0.27 µg/ml.

Target Details

Target:	VEGFR2/CD309 (VEGFR2)
---------	-----------------------

Target Details

Alternative Name: VEGFR2/Flk-1/KDR ([VEGFR2 Products](#))

Background: VEGFR2, also called as KDR or Flk-1, is identified as the receptor for VEGF and VEGFC and an early marker for endothelial cell progenitors, whose expression is restricted to endothelial cells in vivo. VEGFR2 was shown to be the primary signal transducer for angiogenesis and the development of pathological conditions such as cancer and diabetic retinopathy. It has been shown that VEGFR2 is expressed mainly in the endothelial cells, and the expression is upregulated in the tumor vasculature. Thus the inhibition of VEGFR2 activity and its downstream signaling are important targets for the treatment of diseases involving angiogenesis. VEGFR2 transduces the major signals for angiogenesis via its strong tyrosine kinase activity. However, unlike other representative tyrosine kinase receptors, VEGFR2 does not use the Ras pathway as a major downstream signaling but rather uses the phospholipase C-protein kinase C pathway to signal mitogen-activated protein (MAP)-kinase activation and DNA synthesis. VEGFR2 is a direct and major signal transducer for pathological angiogenesis, including cancer and diabetic retinopathy, in cooperation with many other signaling partners; thus, VEGFR2 and its downstream signaling appear to be critical targets for the suppression of these diseases. VEGF and VEGFR2-mediated survival signaling is critical to endothelial cell survival, maintenance of the vasculature and alveolar structure and regeneration of lung tissue. Reduced VEGF and VEGFR2 expression in emphysematous lungs has been linked to increased endothelial cell death and vascular regression. [Immune Checkpoint](#) [Immunotherapy](#) [Cancer Immunotherapy](#) [Targeted Therapy](#)

Synonym: 6130401C07;Flk-1;Flk1;Krd-1;Ly73;sVEGFR-2;VEGFR-2;VEGFR2

Molecular Weight: 110 kDa

UniProt: [P35918](#)

Pathways: [RTK Signaling](#), [Glycosaminoglycan Metabolic Process](#), [Signaling Events mediated by VEGFR1 and VEGFR2](#), [Growth Factor Binding](#), [Regulation of long-term Neuronal Synaptic Plasticity](#), [VEGF Signaling](#)

Application Details

Restrictions: For Research Use only

Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Handling

Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.