

Datasheet for ABIN7320118

Cyclin E1 Protein (CCNE1) (GST tag,His tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	Cyclin E1 (CCNE1)
Origin:	Mouse
Source:	Baculovirus infected Insect Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Cyclin E1 protein is labelled with GST tag,His tag.

Product Details

Purpose:	Recombinant Mouse CCNE1/Cyclin-E1 Protein (His & GST Tag)(Active)
Sequence:	Met1-Glu408
Characteristics:	A DNA sequence encoding the mouse CCNE1 (AAH62152.1) (Met1-Glu408) was expressed with the N-terminal polyhistidine-tagged GST tag at the N-terminus.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized mouse CCNE1 at 10 µg/ml (100 µl/well) can bind biotinylated human CDK4 , The EC50 of biotinylated human CDK4 is 48.0-114.0 ng/ml.

Target Details

Target:	Cyclin E1 (CCNE1)
---------	-------------------

Target Details


Alternative Name:	CCNE1/Cyclin-E1 (CCNE1 Products)
Background:	<p>Background: Cyclin E1 is a member of the highly conserved cyclin family and belongs to the E-type cyclin that functions as a regulator of S phase entry and progression in mammalian cells. Cyclin E1 serves as regulatory subunits that bind, activate, and provide substrate for its associated cyclin-dependent kinase2 (CDK2), whose activity is essential for cell cycle G1 / S transition. Over expression of this encoding gene has been found in many tumors, which results in chromosome instability and by extension, induce tumorigenesis. This protein was also found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. In general, cyclin E1, as an activator of phospho-CDK2 (pCDK2), is important for cell cycle progression and is frequently overexpressed in cancer cells.</p> <p>Synonym: AW538188,CycE1</p>
Molecular Weight:	74.8 kDa
Pathways:	Cell Division Cycle , Intracellular Steroid Hormone Receptor Signaling Pathway , Nuclear Hormone Receptor Binding , Mitotic G1-G1/S Phases

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 50 mM Tris, 100 mM NaCl, 3 mM DTT, 0.5 mM GSH, 10 % glycerol, pH 8.0
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.