antibodies .- online.com







PROCR Protein (Fc Tag)







()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	50 μg
Target:	PROCR
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PROCR protein is labelled with Fc Tag.
Product Details	
Purpose:	Recombinant Mouse EPCR Protein (Fc Tag)
Sequence:	Met 1-Ser 214
Characteristics:	A DNA sequence encoding the mouse PROCR (Q64695) extracellular domain (Met 1-Ser 214) was fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Target Details	
Target:	PROCR
Alternative Name:	EPCR (PROCR Products)
Background:	Background: Endothelial protein C receptor (EPCR), also known as activated protein C receptor

(APC receptor) or PROCR, is a receptor for Protein C. Protein C plays an important role in many metabolism processes in humans and other animals after activated by binding to Endothelial

protein C receptor (EPCR). Because of the EPCR is found primarily on endothelial cells (cells on the inside of blood vessels), activated protein C is found maily near endothelial cells. Protein C is pleiotropic, with two main functions: anticoagulation and cytoprotection. Which function will be performed depend on whether or not protein C remains bind to EPCR after activated. The anticoagulation occurs when it does not. In this case, protein C functions as an anticoagulant by irreversibly proteolytically inactivating Factor Va and Factor VIIIa, turning them into Factor Vi and Factor VIIIi respectively. When still bound to EPCR, activated protein C performs its cytoprotective effects, acting on the effector substrate PAR-1, protease-activated receptor-1. To a degree, APC's anticoagulant properties are independent of its cytoprotective ones, in that expression of one pathway is not affected by the existence of the other.

Synonym: Al325044;Ccca;Ccd41;Epcr

Molecular Weight:

49.3 kDa

UniProt:

Q64695


Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized		
Reconstitution:	Please refer to the printed manual for detailed information.		
Buffer:	Lyophilized from sterile PBS, pH 7.4		
Storage:	4 °C,-20 °C,-80 °C		
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.		
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted		
	samples are stable at < -20°C for 3 months.		



Western Blotting

Image 1.