

Datasheet for ABIN7320140

IL1R1 Protein (Fc Tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	IL1R1
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This IL1R1 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Mouse IL1R1/CD121a Protein (Fc Tag)(Active)
Sequence:	Met 1-Lys 338
Characteristics:	A DNA sequence encoding the mouse IL1R1 (P13504) extracellular domain (Met 1-Lys 338) was fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 92 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	1. Measured by its binding ability in a functional ELISA.2. Immobilized recombinant human IL1b at 10 µg/mL (100 µl/well) can mouse IL1R1-Fc with a linear range of 62.5-1000 ng/mL.

Target Details

Target:	IL1R1
---------	-------

Target Details

Alternative Name:	IL1R1/CD121a (IL1R1 Products)
Background:	<p>Background: Interleukin 1 receptor, type I (IL-1R1) also known as CD121a (Cluster of Differentiation 121a), is an interleukin receptor. IL-1R1/CD121a is a cytokine receptor that belongs to the interleukin 1 receptor family. This protein is a receptor for interleukin alpha (IL1A), interleukin beta (IL1B), and interleukin 1 receptor, type I (IL1R1/IL1RA). IL-1R1/CD121a is an important mediator involved in many cytokine induced immune and inflammatory responses. This protein has been characterized by pharmacological and molecular techniques in the mouse brain. The spindle-shaped astrocytes enclose the wound, separating the healthy from damaged neural tissue. The shape change and subsequent repair processes are IL-1β activity-dependent, acting through the IL-1 type 1 receptor (IL-1R1), as co-application of the IL-1 type 1 receptor antagonist protein (IL-1ra) blocks IL-1β induced effects. In the spleen, a slight increase in IL-1R AcP and IL-1R1 was observed during the first hours following LPS stimulation. In conclusion, IL-1R AcP mRNA is expressed in the brain and in other tissues where IL-1R1/CD121a transcripts are found. However, the regulation of its expression is distinct from IL-1R1/CD121a. The high level of expression and the lack of regulation of IL-1R AcP transcripts in the brain under inflammatory conditions suggest that the protein might be constitutively expressed in excess.</p> <p>Synonym: Interleukin-1 receptor type 1;IL-1RT1; IL-1 RI;CD121b;IL-1R1;IL-iR;Il1r-1</p>
Molecular Weight:	64.3 kDa
UniProt:	P13504
Pathways:	NF-kappaB Signaling , Carbohydrate Homeostasis , Cancer Immune Checkpoints

Application Details

Restrictions:	For Research Use only
---------------	-----------------------


Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

Handling

samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.