

Datasheet for ABIN7320165
CD69 Protein (CD69) (His tag,ECD)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	CD69
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD69 protein is labelled with His tag,ECD.

Product Details

Purpose:	Recombinant Mouse CD69 Protein (His tag, ECD)
Sequence:	Asn62-Arg199
Characteristics:	A DNA sequence encoding the mouse Cd69 (NP_001028294.1) (Asn62-Arg199) was expressed with a polyhistidine tag at the C-terminus.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.

Target Details

Target:	CD69
Alternative Name:	CD69 (CD69 Products)
Background:	Background: Early activation antigen CD69, also known as activation inducer molecule (AIM), is a single-pass type II membrane protein. Recently, cDNA clones encoding human and mouse CD69 were isolated and showed CD69 to be a member of the C-type lectin superfamily. It is one

Target Details

of the earliest cell surface antigens expressed by T cells following activation. Once expressed, CD69 acts as a costimulatory molecule for T cell activation and proliferation. In addition to mature T cells, CD69 is inducibly expressed by immature thymocytes, B cells, natural killer (NK) cells, monocytes, neutrophils and eosinophils, and is constitutively expressed by mature thymocytes and platelets. CD69 is involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets. The structure, chromosomal localization, expression and function of CD69 suggest that it is likely a pleiotropic immune regulator, potentially important in the activation and differentiation of a wide variety of hematopoietic cells. This membrane molecule transiently expresses on activated lymphocytes, and its selective expression in inflammatory infiltrates suggests that it plays a role in the pathogenesis of inflammatory diseases. CD69 plays a crucial role in the pathogenesis of allergen-induced eosinophilic airway inflammation and hyperresponsiveness and that CD69 could be a possible therapeutic target for asthmatic patients.

Synonym: 5830438K24Rik, AI452015, AIM, VEA, CLEC2C

Molecular Weight:	17.3 kDa
-------------------	----------

NCBI Accession:	NP_001028294
-----------------	------------------------------

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling


Format:	Lyophilized
---------	-------------

Reconstitution:	Please refer to the printed manual for detailed information.
-----------------	--

Buffer:	Lyophilized from sterile PBS, pH 7.4
---------	--------------------------------------

Storage:	4 °C, -20 °C, -80 °C
----------	----------------------

Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
------------------	---



Western Blotting

Image 1.