# .-online.com antibodies

## Datasheet for ABIN7320170 SDC4 Protein (AA 1-146) (His tag)

Image



#### Overview

Quantity:	100 µg
Target:	SDC4
Protein Characteristics:	AA 1-146
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SDC4 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Mouse Syndecan-4/SDC4 Protein (aa 1-146, His Tag)
Sequence:	Met 1-Val 146
Characteristics:	A DNA sequence encoding the mouse SDC4 (035988) extracellular domain (Met 1-Val 146) was expressed, fused with a C-terminal polyhistidine tag.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.

#### Target Details

Target:	SDC4
Alternative Name:	Syndecan-4/SDC4 (SDC4 Products)
Background:	Background: SDC4 (Syndecan-4), also known as Syn4, is a transmembrane heparan sulfate


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7320170 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	proteoglycan that co-operates with integrins during cell-matrix interactions for the assembly of
	focal adhesions and actin stress fibers and in the phosphorylation of focal adhesion kinase
	(FAK) on Tyr397. Syndecan-4 plays roles in the formation of focal adhesions and stress fibers.
	The cytoplasmic domain of syndecan-4 interacts with a number of signalling and structural
	proteins, and both extracellular and cytoplasmic domains are necessary for regulated activation
	of associated transmembrane receptors. Syndecan-4/SDC4 is a heparan sulfate proteoglycan
	and works as a coreceptor for various growth factors. SDC4 deficiency limits neointimal
	formation after vascular injury by regulating vascular smooth muscle cells (VSMCs)
	proliferation and vascular progenitor cells (VPCs) mobilization. Therefore, SDC4 may be a novel
	therapeutic target for preventing arterial restenosis after angioplasty.
	Synonym: AA959608,AW108331,ryudocan,Synd4,syndecan-4,SDC4,SYND4,Ryudocan core
	protein
Molecular Weight:	14.9 kDa
UniProt:	035988
Pathways:	Glycosaminoglycan Metabolic Process

## Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7320170 | 09/09/2023 | Copyright antibodies-online. All rights reserved.



Western Blotting

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7320170 | 09/09/2023 | Copyright antibodies-online. All rights reserved.