

Datasheet for ABIN7320178

CD45 Protein (Fc Tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	CD45 (PTPRC)
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD45 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Mouse CD45/PTPRC Protein (Fc Tag)(Active)
Sequence:	Met1-Lys425
Characteristics:	A DNA sequence encoding the mouse PTPRC (NP_035340.3) (Met1-Lys425) was expressed with the Fc region of human IgG1 at the C-terminus.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized human SEMA4D-His at 10 µg/ml (100 µl/well) can bind mouse PTPRC-Fc with a linear range of 0.625-5 µg/ml.

Target Details

Target:	CD45 (PTPRC)
---------	--------------

Target Details

Alternative Name:	CD45/PTPRC (PTPRC Products)
Background:	<p>Background: The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Protein tyrosine phosphatase, receptor type C (CD45), also known as PTPRC is a member of the protein tyrosine phosphatase (PTP) family which is known for its function to serve as signaling molecules and to regulate a variety of cellular processes such as cell proliferation, differentiation, mitotic cycle and oncogenic transformation. CD45 is found expression specifically in hemotopietic cells. CD45 consists of an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains. It serves as an essential regulator of T-cell and B-cell antigen receptor signaling through either direct interaction with components of the antigen receptor complexes or by activating various Src family kinases required for the antigen receptor signaling and it also can suppress JAK kinases.</p> <p>Synonym: B220,Cd45,CD45R,L-CA,loc,Ly-5,Lyt-4,T200</p>
Molecular Weight:	71.7 kDa
NCBI Accession:	NP_035340
Pathways:	TCR Signaling , Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Production of Molecular Mediator of Immune Response , CXCR4-mediated Signaling Events , BCR Signaling


Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C

Handling

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.