antibodies .- online.com





FZD10 Protein (His tag)





Go to Product page

\sim					
	1//	r۱.	/ I	\triangle	٨

Quantity:	100 μg
Target:	FZD10
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FZD10 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse Frizzled-10/FZD10 Protein (His Tag)	
Sequence:	Met 1-Gly 162	
Characteristics:	A DNA sequence encoding the N-terminal segment of mouse FZD10 (NP_780493.1) (Met 1-Gly 162) was expressed, fused with a polyhistidine tag at the C-terminus.	
Purity:	> 95 % as determined by SDS-PAGE	
Endotoxin Level:	$<$ 1.0 EU per μg of the protein as determined by the LAL method.	

Target Details

Target:	FZD10	
Alternative Name:	Frizzled-10/FZD10 (FZD10 Products)	
Background:	Background: Frizzled-10, also known as Fz-10, CD350 and FZD10, is a multi-pass membrane protein which belongs to the G-protein coupled receptor Fz/Smo family. Frizzled-10 / FZD10 abundantly expressed in the cerebellum, followed by cerebral cortex, medulla and spinal cortex.	

very low levels in total brain, frontal lobe, temporal lobe and putamen. It is weakly expressed in adult brain, heart, lung, skeletal muscle, pancreas, spleen and prostate. Frizzled-10 / FZD10 is a receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. Frizzled-10 / FZD10 may also be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.

Synonym: Fz-10

Molecular Weight: 17.4 kDa

NCBI Accession: NP_780493


Pathways: WNT Signaling

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.