

Datasheet for ABIN7320240

**DDR2 Protein (His tag)****1** Image[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	DDR2
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This DDR2 protein is labelled with His tag.

## Product Details

Purpose:	Recombinant Mouse DDR2 Kinase/CD167b Protein (His Tag)(Active)
Sequence:	Met 1-Arg 399
Characteristics:	A DNA sequence encoding the extracellular domain of mouse DDR2 (NP_072075.2) (Met 1-Arg 399) was expressed, with a polyhistidine tag at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to bind rat tail Collagen I in a functional ELISA.

## Target Details

Target:	DDR2
Alternative Name:	DDR2 ( <a href="#">DDR2 Products</a> )

## Target Details

Background:	<p>Background: Discoidin domain receptor 2 (DDR2) or CD167b (cluster of differentiation 167b) is a kind of protein tyrosine kinases associated with cell proliferation and tumor metastasis, and collagen, identified as a ligand for DDR2, up-regulates matrix metalloproteinase 1 (MMP-1) and MMP-2 expression in cellular matrix. DDR2/CD167b was found to recognise the triple-helical region of collagen X as well as the NC1 domain. Binding to the collagenous region was dependent on the triple-helical conformation. DDR2/CD167b autophosphorylation was induced by the collagen X triple-helical region but not the NC1 domain, indicating that the triple-helical region of collagen X contains a specific DDR2 binding site that is capable of receptor activation. DDR2/CD167b is induced during stellate cell activation and implicate the phosphorylated receptor as a mediator of MMP-2 release and growth stimulation in response to type I collagen. Moreover, type I collagen-dependent upregulation of DDR2/CD167b expression establishes a positive feedback loop in activated stellate cells, leading to further proliferation and enhanced invasive activity.</p> <p>Immune Checkpoint   Immunotherapy   Cancer Immunotherapy   Targeted Therapy</p> <p>Synonym: Discoidin domain-containing receptor 2; Discoidin domain receptor 2; CD167 antigen-like family member B; Neurotrophic tyrosine kinase; receptor-related 3; Receptor protein-tyrosine kinase TKT; Tyrosine-protein kinase TYRO10; CD167b; Ddr2;Ntrkr3;tyro10</p>
-------------	--

Molecular Weight:	44 kDa
-------------------	--------

NCBI Accession:	<a href="#">NP_072075</a>
-----------------	---------------------------

Pathways:	<a href="#">RTK Signaling</a>
-----------	-------------------------------

## Application Details

Comment:	80-90 kDa
----------	-----------

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Lyophilized
---------	-------------

Reconstitution:	Please refer to the printed manual for detailed information.
-----------------	--

Buffer:	Lyophilized from sterile PBS, pH 7.4
---------	--------------------------------------


Storage:	4 °C,-20 °C,-80 °C
----------	--------------------

Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
------------------	---

Handling

samples are stable at < -20°C for 3 months.

Images



**Western Blotting**

**Image 1.**