.-online.com antibodies

Datasheet for ABIN7320241 CD2 Protein (CD2) (His tag)

Image



Overview

Quantity:	50 µg
Target:	CD2
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD2 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse CD2 Protein (His Tag)
Sequence:	Met 1-Ser 203
Characteristics:	A DNA sequence encoding the extracellular domain of mouse CD2 (NP_038514.1) (Met 1-Ser 203) was expressed, with a polyhistidine tag at the C-terminus.
Purity:	> 98 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per μ g of the protein as determined by the LAL method.


Target Details

Target:	CD2
Alternative Name:	CD2 (CD2 Products)
Background:	Background: T-cell surface antigen CD2, also known as T-cell surface antigen T11/Leu-5, and SRBC, is a single-pass type I membrane protein. It contains one Ig-like C2-type domain and one
	Ig-like V-type domain. CD2 is a cell adhesion molecule expressed on T cells and is recognized

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7320241 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	as a target for CD48 (rats) and CD58 (humans). CD2 has been shown to set quantitative
	thresholds in T cell activation both in vivo and in vitro. Further, intracellular CD2 signaling
	pathways and networks are being discovered by the identification of several cytosolic tail
	binding proteins. CD2 interacts with lymphocyte function-associated antigen (LFA-3) and
	CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the
	triggering of T-cells, the cytoplasmic domain of CD2 is implicated in the signaling function. The
	complex of CD2 and CD58 also plays an important role in enhancing the adhesion of T
	lymphocytes to target cells, and promoting hyperplasia and activation of T lymphocytes. As a
	cell surface glycoprotein, CD2 expressed on most human T cells and natural killer (NK) cells
	and plays an important role in mediating cell adhesion in both T-lymphocytes and in signal
	transduction.Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted
	Therapy
	Synonym: LFA-2;Ly-37;Ly37
Molecular Weight:	22 kDa
NCBI Accession:	NP_038514
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7320241 | 09/09/2023 | Copyright antibodies-online. All rights reserved.



Western B	lotting
-----------	---------

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7320241 | 09/09/2023 | Copyright antibodies-online. All rights reserved.