.-online.com antibodies

Datasheet for ABIN7320251 CD99 Protein (CD99) (Fc Tag)

Image



Overview

Quantity:	100 µg
Target:	CD99
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD99 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Mouse CD99/PILR1 Protein (Fc Tag)	
Sequence:	Met 1-Gly 137	
Characteristics:	A DNA sequence encoding the mouse CD99 (NP_612182.1) extracellular domain (Met 1-Gly 137) was fused with the Fc region of human IgG1 at the C-terminus.	
Purity:	> 90 % as determined by SDS-PAGE	
Endotoxin Level:	< 1.0 EU per μ g of the protein as determined by the LAL method.	

Target Details

Target:	CD99
Alternative Name:	CD99/PILR1 (CD99 Products)
Target Type:	Viral Protein
Background:	Background: The cluster of differentiation (CD) system is commonly used as cell markers in

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7320251 | 09/09/2023 | Copyright antibodies-online. All rights reserved. immunophynotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD99 is a transmembrane protein expressed on most hematopoietic cells, endothelial cells and at the borders between confluent cells. CD99 is also found expressed in the development of normal ovary and testis as well as in 25 sex cord-stromal tumors, 7 epithelial neoplasms, and 6 germ cell tumors. CD99 may be a useful marker for sex cord-stromal tumors and that its degree of reactivity correlates with the degree of differentiation in Sertoli-Leydig cell tumors. Additionally, CD99 might aid in distinguishing granulose cell tumors of the ovary from poorly differentiated carcinomas and it has been reported to be a sensitive and specific marker for Ewing's sarcoma and primitive neuroectodermal tumor.

Synonym: 1110061M03Rik,2410026K10Rik,D4,pilr-1,Pilr-I

Molecular Weight:	37.5 kDa
NCBI Accession:	NP_612182

Application Details


Restrictions:

For Research Use only

Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7320251 | 09/09/2023 | Copyright antibodies-online. All rights reserved.



Western B	lotting
-----------	---------

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7320251 | 09/09/2023 | Copyright antibodies-online. All rights reserved.