.-online.com antibodies

Datasheet for ABIN7320277 ERK2 Protein (GST tag,His tag)

Image



Overview

Quantity:	50 µg
Target:	ERK2 (MAPK1)
Origin:	Mouse
Source:	Baculovirus infected Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ERK2 protein is labelled with GST tag,His tag.

Product Details

Purpose:	Recombinant Mouse ERK2/MAPK1/MAPK2 Protein (His & GST Tag)
Sequence:	Met1-Ser358
Characteristics:	A DNA sequence encoding the mouse MAPK1 (P63085) (Met1-Ser358) was fused with the N- terminal polyhistidine-tagged GST tag at the N-terminus.
Purity:	> 96 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per μ g of the protein as determined by the LAL method.

Target Details

Target:	ERK2 (MAPK1)
Alternative Name:	ERK2/MAPK1/MAPK2 (MAPK1 Products)
Background:	Background: MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular
	processes such as proliferation, differentiation, transcription regulation and development. ERK is a

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7320277 | 09/09/2023 | Copyright antibodies-online. All rights reserved. versatile protein kinase that regulates many cellular functions. Growing evidence suggests that extracellular signal-regulated protein kinase 1/2 (ERK1/2) plays a crucial role in promoting cell death in a variety of neuronal systems, including neurodegenerative diseases. It is believed that the magnitude and the duration of ERK1/2 activity determine its cellular function. Activation of ERK1/2 are implicated in the pathophysiology of spinal cord injury (SCI). ERK2 signaling is a novel target associated with the deleterious consequences of spinal injury. ERK-2, also known as Mitogenactivated protein kinase 1 (MAPK1), is a member of the protein kinase superfamily and MAP kinase subfamily. MKP-3 is a dual specificity phosphatase exclusively specific to MAPK1 for its substrate recognition and dephosphorylating activity. The activation of MAPK1 requires its phosphorylation by upstream kinases. Upon activation, MAPK1 translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. MAPK1 is involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK1. MAPK1 acts as a transcriptional repressor which represses the expression of interferon gamma-induced genes. Transcriptional activity is independent of kinase activity. The nuclear-cytoplasmic distribution of ERK2 is regulated in response to various stimuli and changes in cell context. Furthermore, the nuclear flux of ERK2 occurs by several energy- and carrier-dependent and -independent mechanisms. ERK2 has been shown to translocate into and out of the nucleus by facilitated diffusion through the nuclear pore, interacting directly with proteins within the nuclear pore complex, as well as by karyopherinmediated transport. ERK2 interacts with the PDE4 catalytic unit by binding to a KIM (kinase interaction motif) docking site located on an exposed beta-hairpin loop and an FQF (Phe-GIn-Phe) specificity site located on an exposed alpha-helix. These flank a site that allows phosphorylation by ERK, the functional outcome of which is orchestrated by the N-terminal UCR1/2 (upstream conserved region 1 and 2) modules.Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy Synonym:

9030612K14Rik;AA407128;AU018647;C78273;ERK;Erk2;MAPK2;p41mapk;p42mapk;Prkm1;PRKM2

Molecular Weight:	69.1 kDa
UniProt:	P63085
Pathways:	MAPK Signaling, RTK Signaling, Apoptosis, Interferon-gamma Pathway, Fc-epsilon Receptor
	Signaling Pathway, Response to Growth Hormone Stimulus, Activation of Innate immune
	Response, Cellular Response to Molecule of Bacterial Origin, Hepatitis C, Protein targeting to
	Nucleus, Toll-Like Receptors Cascades, Monocarboxylic Acid Catabolic Process, Autophagy, G-
	protein mediated Events, Signaling Events mediated by VEGFR1 and VEGFR2, Signaling of
	Hepatocyte Growth Factor Receptor, VEGFR1 Specific Signals, BCR Signaling, S100 Proteins

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7320277 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Application Details


Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 20 mM Tris, 500 mM NaCl, pH 8.0, 10 % glycerol
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 $^{\circ}$ C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7320277 | 09/09/2023 | Copyright antibodies-online. All rights reserved.