# antibodies - online.com







# CD147 Protein (AA 1-209) (His tag)



# Image



### Overview

Quantity:	100 μg
Target:	CD147 (BSG)
Protein Characteristics:	AA 1-209
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD147 protein is labelled with His tag.

## **Product Details**

Purpose:	Recombinant Mouse CD147/Basigin Protein (aa 1-209, His Tag)
Sequence:	Met 1-Arg 209
Characteristics:	A DNA sequence encoding the extracellular domain of mouse BSG isoform 2 (NP_001070652.1) (Met 1-Arg 209) was expressed, fused with a polyhistidine tag at the C-terminus.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
T . D . "	

# **Target Details**

Target:	CD147 (BSG)
Alternative Name:	CD147/Basigin (BSG Products)

### Target Details

Ranko	round:
Dacke	n Ouriu.

Background: CD147/EMMPRIN (Extracellular Matrix Metalloproteinase Inducer), also known as Basigin (BSG), is a transmembrane glycoprotein with different forms resulted from different modes of glycosylation and N-terminal sequence variants. It is a member of the immunoglobulin superfamily with homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. This protein play important roles in variety of events including spermatogenesis, embryo implantation, neural network formation. CD147 induces the production and release of matrix metalloproteinases (MMP) in the surrounding mesenchymal cells and tumor cells, and thereby promotes invasion, metastasis, growth and survival of malignant cells. Furthermore, CD147 also serves as a receptor for extracellular cyclophilinthe and its association with integrins might be important in signal transduction. Recently, CD147 displays increased expression in many cancers, and it has been previously demonstrated to participate in cancer metastasis and progression. Thus, CD147 and its antibody are used as an effective treatment for malignant cancers.

Synonym: Basigin, Basic immunoglobulin superfamily, HT7 antigen, Membrane glycoprotein

Synonym: Basigin, Basic immunoglobulin superfamily, HT7 antigen, Membrane glycoprotein gp42, CD147, Bsg,Al115436,Al325119,EMMPRIN,HT-7

Mo	lacular	Weiaht:
IVIO	ıccuiai	vveigiii.

30 kDa

NCBI Accession:

NP\_001070652

Pathways:

S100 Proteins


### **Application Details**

Restrictions:

For Research Use only

### Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	



# **Western Blotting**

Image 1.