

Datasheet for ABIN7320338
HDAC8 Protein (His tag)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	HDAC8
Origin:	Mouse
Source:	Baculovirus infected Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HDAC8 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse HDAC8/HDACL1 Protein (His Tag)
Sequence:	Met 1-Val 377
Characteristics:	A DNA sequence encoding the mouse HDAC8 isoform 1 (Q8VH37-1) (Met 1-Val 377) was expressed, with a C-terminal polyhistidine tag.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.

Target Details

Target:	HDAC8
Alternative Name:	HDAC8/HDACL1 (HDAC8 Products)
Background:	Background: Histone deacetylase 8, also known as HDAC8 and HDACL1, is a nucleus and cytoplasm protein which belongs to the histone deacetylase family and HD type 1 subfamily. Histone deacetylases (HDACs) are a growing family of enzymes implicated in transcriptional

Target Details

regulation by affecting the acetylation state of core histones in the nucleus of cells. HDAC8 / HDACL1 is weakly expressed in most tissues. It expressed at higher level in heart, brain, kidney and pancreas and also in liver, lung, placenta, prostate and kidney. HDAC8 / HDACL1 is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. HDAC8 / HDACL1 may play a role in smooth muscle cell contractility. HDAC8 / HDACL1 may be a potential drug target for neuroblastoma differentiation therapy using selective inhibitors, avoiding unspecific side effects.

Synonym: 2610007D20Rik

Molecular Weight: 43.1 kDa

Pathways: [Cellular Glucan Metabolic Process](#)

Application Details

Restrictions: For Research Use only

Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile 20 mM Tris, 500 mM NaCl, pH 7.4, 10 % glycerol

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.