

Datasheet for ABIN7320346

Podoplanin Protein (PDPN) (His tag,Fc Tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	Podoplanin (PDPN)
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Podoplanin protein is labelled with His tag,Fc Tag.

Product Details

Purpose:	Recombinant Mouse Podoplanin/PDPN Protein (His & Fc Tag)(Active)
Sequence:	Met 1-Leu 141
Characteristics:	A DNA sequence encoding the extracellular domain (Met 1-Leu 141) of mouse PDPN (NP_034459.2) precursor was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Purity:	> 97 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Immobilized mouse PDPN-Fch at 10 µg/ml (100 µl/well) can bind biotinylated human CLEC1B-His, The EC50 of biotinylated human CLEC1B-His is 0.04-0.08 µg/ml.

Target Details

Target:	Podoplanin (PDPN)
---------	-------------------

Target Details


Alternative Name:	Podoplanin/PDPN (PDPN Products)
Background:	<p>Background: Podoplanin, also known as PDPN, is a type-I integral membrane glycoprotein with diverse distribution in human tissues. The physiological function of this protein may be related to its mucin-type character. The homologous protein in other species has been described as a differentiation antigen and influenza-virus receptor. The specific function of this protein has not been determined. Alternatively spliced transcript variants encoding different isoforms have been identified. PDPN is a mucin-type glycoprotein negatively charged by extensive O-glycosylation and a high content of sialic acid, which expresses the adhesive property. It is selectively expressed in lymphatic endothelium as well as lymphangiomas, Kaposi sarcomas, and in a subset of angiosarcomas with probable lymphatic differentiation. PDPN may contribute to form odontoblastic fiber or function as the anchorage to the tooth development and in proliferating epithelial cells of cervical loop and apical bud. The intensity of podoplanin expression is negatively correlated with the expression of CD34 and factor VIII. Podoplanin would be useful as a diagnostic marker for epithelioid hemangioendothelioma in liver tumors.</p> <p>Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy</p> <p>Synonym: Gp38;OTS-8;RANDAM-2;T1-alpha;T1a;T1alpha;Podoplanin; Aggrus; Glycoprotein 38; PA2.26 antigen</p>
Molecular Weight:	40.6 kDa
NCBI Accession:	NP_034459
Pathways:	Dicarboxylic Acid Transport

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C, -20 °C, -80 °C
Storage Comment:	<p>Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.</p> <p>Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>



Western Blotting

Image 1.