.-online.com antibodies

Datasheet for ABIN7320353 **PEDF Protein (His tag)**

Image



Overview

Quantity:	50 µg
Target:	PEDF (SERPINF1)
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This PEDF protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse Serpin F1/PEDF Protein (His Tag)(Active)
Sequence:	Met 1-Thr 417
Characteristics:	A DNA sequence encoding the extracellular domain (Met 1-Thr 417) of mouse SerpinF1 (NP_035470.3) precursor was expressed with a C-terminal polyhistidine tag.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per μ g of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA.Immobilized mouse SERPINF1-His at 10 μ g/ml (100 μ l/well) can bind biotinylated human GST-CSNK2A1 with a linear range of 0.31-2.5 μ g/ml.

Target Details

Target:

PEDF (SERPINF1)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7320353 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	Serpin F1/PEDF (SERPINF1 Products)
Background:	Background: Pigment epithelium-derived factor, also known as PEDF, Serpin F1, and SERPINF1,
	is a multiple functional protein which has both anti-angiogenic activity and neurotrophic activity
	at the same time. PEDF is a secreted glycoprotein that belongs to the noninhibitory serpin. It
	has an alpha/beta core serine-protease inhibitor domain, three major beta-sheets, and ten
	alpha-helices. PEDF does not inhibit either serine or cysteine proteinases. PEDF exerts diverse
	physiological activities including anti-angiogenesis, anti-vasopermeability, anti-tumor, and
	neurotrophic activities. PEDF acts via multiple high affinity ligands and cell receptors. It has
	been described as a natural angiogenesis inhibitor with neurotrophic and immune-modulation
	properties. PEDF induces macrophages apoptosis and necrosis through the activation of
	peroxisome proliferator-activated receptor-gamma by which PEDF could modulate
	inflammatory reactions in septic shock. It balances angiogenesis in the eye and blocks tumor
	progression.
	Synonym: Al195227 Protein, Mouse,EPC-1 Protein, Mouse,Pedf Protein, Mouse,Pedfl Protein,
	Mouse,Sdf3 Protein, Mouse
Molecular Weight:	45.8 kDa
NCBI Accession:	NP_035470
Application Details	


Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7320353 | 09/09/2023 | Copyright antibodies-online. All rights reserved.



Western Blotting

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7320353 | 09/09/2023 | Copyright antibodies-online. All rights reserved.