

Datasheet for ABIN7320411

IL7R Protein (His tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	IL7R
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL7R protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse IL7RA/CD127 Protein (His Tag)
Sequence:	Met 1-Asp 239
Characteristics:	A DNA sequence encoding the mouse IL7Rα (NP_032398.3) extracellular domain (Met 1-Asp 239) was fused with a polyhistidine tag at the C-terminus.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.

Target Details

Target:	IL7R
Alternative Name:	IL7RA/CD127 (IL7R Products)
Background:	Background: Interleukin 7 Receptor alpha (IL-7RA), also known as CD127, is a 75 kDa hematopoietin receptor superfamily member that plays an important role in lymphocyte differentiation, proliferation, and survival. IL-7 receptor alpha (CD127) signaling is essential for

Target Details

T-cell development and regulation of naive and memory T-cell homeostasis. IL-7RA is critically required for the proper development and function of lymphoid cells. Therefore, the IL-7RA is critically required for the proper development and function of lymphoid cells. Studies from both pathogenic and controlled HIV infection indicate that the containment of immune activation and preservation of CD127 expression are critical to the stability of CD4(+) T cells in infection. A better understanding of the factors regulating CD127 expression in HIV disease, particularly on T(CM) cells, might unveil new approaches exploiting the IL-7/IL-7R receptor pathway to restore T cell homeostasis and promote immune reconstitution in HIV infection. Factors relevant to HIV infection that could potentially decrease CD127 expression on human CD8(+) T cells. CD127 down-regulation may be an important contributor to HIV-associated T-cell dysfunction. In addition to IL-7, IL-7RA also associates with TSLPR to form the functional receptor for thymic stromal lymphopoietin (TSLP) which indirectly regulates T cell development by modulating dendritic cell activation. Mutations in the human IL-7RA gene cause a type of severe combined immunodeficiency in which the major deficiencies are in T cell development, whereas B and NK cells are relatively normal in number. Variation in the IL7RA gene was recently found associated with multiple sclerosis (MS). The polymorphisms in the IL7RA gene is involved in MS pathogenesis and suggest that IL7RA variation may primarily affect chronic disease courses. Soluble CD127 (sCD127) appears to play an important role in the immunopathogenesis of several chronic infections, multiple sclerosis, and various cancers.

Synonym: Interleukin-7 receptor subunit alpha, IL7r, IL-7 receptor subunit alpha, IL-7R subunit alpha, IL-7R-alpha, IL-7RA, CD127,IL-7Ralpha

Molecular Weight:	26.5 kDa
NCBI Accession:	NP_032398
Pathways:	JAK-STAT Signaling , Regulation of Leukocyte Mediated Immunity , Production of Molecular Mediator of Immune Response , Regulation of Cell Size


Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4

Handling

Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.