

Datasheet for ABIN7320423

Follistatin Protein (FST) (His tag)[Go to Product page](#)**1** Image

Overview

Quantity:	50 µg
Target:	Follistatin (FST)
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Follistatin protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse Follistatin/FST Protein (His Tag)(Active)
Sequence:	Met 1-Trp 344
Characteristics:	A DNA sequence encoding the mouse FST (P47931) (Met 1-Trp 344) was expressed, fused with a polyhistidine tag at the C-terminus.
Purity:	> 98 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED50 for this effect is typically 0.02-0.08 µg/mL in the presence of 10 ng/ml Recombinant Human Activin A.

Target Details


Target:	Follistatin (FST)
---------	-------------------

Target Details

Alternative Name:	Follistatin/FST (FST Products)
Background:	<p>Background: Follistatin is a single-chain gonadal protein that specifically inhibits follicle-stimulating hormone release. The single FST gene encodes two isoforms, FST317 and FST344 containing 317 and 344 amino acids respectively, resulting from alternative splicing of the precursor mRNA. In a study in which 37 candidate genes were tested for linkage and association with polycystic ovary syndrome (PCOS) or hyperandrogenemia in 150 families, evidence was found for linkage between PCOS and follistatin. follistatin are expressed and subserve local regulatory roles in numerous extragonadal tissues, including brain, adrenal, bone marrow, and placenta but perhaps most notably in anterior pituitary-the classical target tissue for inhibin, the activin-follistatin system may play a key role in early embryogenesis. Follistatin binds directly to activin and functions as an activin antagonist. Specific inhibitor of the biosynthesis and secretion of pituitary follicle stimulating hormone follistatin is a binding protein to activin. Since activin binds to follistatin, it is imperative to determine the nature of the activin/follistatin binding complex.</p> <p>Synonym: AL033346,FS</p>
Molecular Weight:	36.1 kDa
UniProt:	P47931
Pathways:	Negative Regulation of Hormone Secretion

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	<p>Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>



Western Blotting

Image 1.