

Datasheet for ABIN7320454
FGFR3 Protein (His tag,Fc Tag)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	FGFR3
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This FGFR3 protein is labelled with His tag,Fc Tag.

Product Details

Purpose:	Recombinant Mouse FGFR3 Protein (His & Fc Tag)(Active)
Sequence:	Met 1-Tyr 367
Characteristics:	A DNA sequence encoding the extracellular domain (Met 1-Tyr 367) of mouse FGFR3 (NP_032036.2) precursor was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Purity:	> 92 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to bind mouse aFGF in a functional ELISA.

Target Details

Target:	FGFR3
---------	-------

Target Details

Alternative Name:	FGFR3 (FGFR3 Products)
Background:	<p>Background: FGFR3, also known as CD333, is a member of the fibroblast growth factor receptor (FGFR) family, with its amino acid sequence being highly conserved between members and among divergent species. FGFR family members differ from one another in their ligand affinities and tissue distribution. FGFRs are transmembrane catalytic receptors that have intracellular tyrosine kinase activity. Mutations in FGFR genes are the cause of several human developmental disorders characterized by skeletal abnormalities such as achondroplasia, and upregulation of FGFR expression may lead to cell transformation and cancer. FGFR3, a full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of FGFR3 interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. FGFR3 binds acidic and basic fibroblast growth hormone and plays a role in bone development and maintenance. Mutations in FGFR3 gene lead to craniosynostosis and multiple types of skeletal dysplasia. Three alternatively spliced transcript variants that encode different protein isoforms have been described. CD333 is the receptor for acidic and basic fibroblast growth factors.</p> <p>Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy</p> <p>Synonym: CD333;Fgfr-3;Flg-2;FR3;HBGFR;Mfr3;sam3</p>
Molecular Weight:	66 kDa
NCBI Accession:	NP_032036
Pathways:	RTK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Stem Cell Maintenance , Growth Factor Binding


Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C, -20 °C, -80 °C

Handling

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.