.-online.com antibodies

Datasheet for ABIN7320659 CRLF2 Protein (Fc Tag)

Image



Overview

Quantity:	50 µg
Target:	CRLF2
Origin:	Mouse
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CRLF2 protein is labelled with Fc Tag.

Product Details


Purpose:	Recombinant Mouse TSLP Receptor/CRLF2 Protein (Fc Tag)
Sequence:	Ala20-Leu233
Characteristics:	Recombinant Mouse Thymic stromal lymphopoietin protein receptor is produced by our Mammalian expression system and the target gene encoding Ala20-Leu233 is expressed with a Fc tag at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	CRLF2
Alternative Name:	TSLP Receptor/CRLF2 (CRLF2 Products)
Background:	Background: The cytokine thymic stromal lymphopoietin receptor (TSLPR) is consisting of a
	common γ receptor-like chain (TSLPR- γ) and a common interleukin 7 (IL-7) R α chain that

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7320659 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	belongs to the type 1 cytokine receptor family. Transfection of TSLPR cDNA result in only low			
	affinity binding, while cotransfection of the IL-7R $lpha$ chain cDNA shows high affinity binding.			
	TSLP and TSLPR play a critical role in the initiation of allergic diseases in mice. The TSLP R cDNA encodes a transmembrane receptor containing 370 amino acids (aa) with two potential			
	N-linked glycosylation sites and a cytoplasmic domain of 104 aa including a single tyrosine			
	residue. TSLPR can mediate signaling of the signal transducer and activator of transcription 5			
	(Stat5) by TSLP. TSLP R is broadly expressed in the immune and hematopoietic cells,			
	particularly in hematopoietic progenitors and myeloid cells.			
	Synonym: CRL2, CRLF2,CRL2 cytokine receptor,cytokine receptor-like factor 2, ILXR, IL-XR,			
	P2RY8/CRLF2 fusion, Thymic stromal lymphopoietin protein receptor, Thymic stromal-derived			
	lymphopoietin receptor, TSLP receptor, TSLPR			
Molecular Weight:	49.8 kDa			
UniProt:	Q8CII9			
Application Details				
Restrictions:	For Research Use only			
Handling				
Format:	Lyophilized			
Reconstitution:	Please refer to the printed manual for detailed information.			
Buffer:	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.			
Storage:	4 °C,-20 °C,-80 °C			
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.			
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted			
	samples are stable at < -20°C for 3 months.			



Western	B	lotting	
---------	---	---------	--

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7320659 | 09/09/2023 | Copyright antibodies-online. All rights reserved.