antibodies -online.com





SNCA Protein (His tag)





Overview

Quantity:	50 μg
Target:	SNCA
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SNCA protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse α-Synuclein/SNCA Protein (His Tag)
Sequence:	Met1-Ala140
Characteristics:	Recombinant Mouse alpha-Synuclein is produced by our E.coli expression system and the target gene encoding Met1-Ala140 is expressed with a 6His tag at the N-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	SNCA
Alternative Name:	alpha-Synuclein/SNCA (SNCA Products)
Background:	Background: Alpha-synuclein (Snca) belongs to a family of proteins including a-, b-, and g-synucleins. Alpha-synuclein has been found to be implicated in the pathophysiology of many
	neurodegenerative diseases, including Parkinson's disease (PD) and Alzheimer's disease.

Manyneurodegenerative diseases has shown that alpha-synuclein accumulates in dystrophic neurites and in Lewy bodies. The function of alpha-synuclein is closely correlated with its three-dimensional structure, especially for proteins important in the pathogenesis of neurodegenerative diseases. Alpha-synuclein is a dynamic molecule whose secondary structure depends on the environment. For example, it has an unfolded random coil structure in aqueous solution, forms a-helical structure upon binding to acidic phospholipid vesicles, and forms insoluble fibrils with a high b-sheet content that resemble the filaments found in Lewy bodies. Also, alpha-synuclein was known to associate with 14-3-3 proteins including protein kinase C, BAD, and extracellular regulated kinase, and overexpression of alpha-synuclein could contribute to cell death in neurodegenerative diseases.

Synonym: Alpha-synuclein, Non-A beta component of AD amyloid, Non-A4 component of amyloid precursor, NACP, Snca

Molecular Weight: 15.9 kDa

055042

Pathways:

UniProt:


Synaptic Membrane, Regulation of G-Protein Coupled Receptor Protein Signaling, Positive
Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process, Plateletderived growth Factor Receptor Signaling, Negative Regulation of Transporter Activity,
Regulation of long-term Neuronal Synaptic Plasticity

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.