antibodies -online.com







TIM3 Protein (AA 20-193) (His tag)





	Go to	Prod	uct	page
--	-------	------	-----	------

_					
U	V	er	VI	е	W

Quantity:	50 μg
Target:	TIM3 (TIM 3)
Protein Characteristics:	AA 20-193
Origin:	Mouse
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TIM3 protein is labelled with His tag.

Product Details


Purpose:	Recombinant Mouse TIM-3/HAVCR2 Protein (aa 20-193, His Tag)
Sequence:	Arg20-Ala193
Characteristics:	Recombinant Mouse T cell Immunoglobulin and Mucin Domain-3 is produced by our Mammalian expression system and the target gene encoding Arg20-Ala193 is expressed with a 6His tag at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	TIM3 (TIM 3)
Alternative Name:	TIM-3/HAVCR2 (TIM 3 Products)

Target Details

Target Type:	Virus
Background:	Background: T cell immunoglobulin and mucin domain-3 (TIM3), also called hepatitis A virus
	cellular receptor 2 (HAVCR2), is a transmembrane glycoprotein of the TIM family of immune
	regulating molecules and plays an important role in the Th1-mediated immune response. TIM3
	is expressed on the Th1 cells, CD8 T-cells, monocytes, and dendritic cells, but not on Th2 cells.
	TIM3 expressed by monocytes and dendritic cells facilitates phagocytosis of apoptotic cells
	and up-regulates cross-presentation of apoptotic cell-associated antigens through interaction
	with phosphatidylserine. Engagement of TIM3 by its ligand galectin-9 induces a range of
	immunosuppressive functions which enhance immune tolerance and inhibit anti-tumor
	immunity. Stimulation of TIM3 with an agonistic antibody promotes inflammation through the
	activation of innate immune cells. TIM3 is also regarded as a potential target molecule for
	immunotherapy. TIM3 and programmed cell death 1 (PD-1) as two important coinhibitory
	regulators of T cell responses, have been implicated with the T-cell dysfunction or exhaustion
	associated with chronic HBV infection including HBV-related HCC.
	Synonym: Hepatitis A virus cellular receptor 2 homolog,HAVcr-2,T-cell immunoglobulin and
	mucin domain-containing protein 3,T-cell immunoglobulin mucin receptor 3,T-cell membrane
	protein 3,Tim3, Timd3
Molecular Weight:	19.9 kDa
Pathways:	Regulation of Lipid Metabolism by PPARalpha, Cancer Immune Checkpoints
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.