

Datasheet for ABIN7320774

**TIM3 Protein (Fc Tag)****1** Image[Go to Product page](#)

## Overview

Quantity:	50 µg
Target:	TIM3 (TIM 3)
Origin:	Mouse
Source:	Human Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This TIM3 protein is labelled with Fc Tag.

## Product Details

Purpose:	Recombinant Mouse TIM-3/HAVCR2 Protein (Fc Tag)(Active)
Sequence:	Arg20-Arg191
Characteristics:	Recombinant Mouse TIM-3 is produced by our Mammalian expression system and the target gene encoding Arg20-Arg191 is expressed with a Fc tag at the C-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Immobilized Galectin 9-His(Cat: PKSH032470) at 2µg/ml(100 µl/well) can bind Mouse TIM-Fc. The ED50 of Mouse TIM-Fc is 4.053ug/ml .

## Target Details

Target:	TIM3 (TIM 3)
---------	--------------

## Target Details

Alternative Name:	TIM-3/HAVCR2 ( <a href="#">TIM 3 Products</a> )
Target Type:	Virus
Background:	<p>Background: T cell immunoglobulin and mucin domain-3 (TIM3), also called hepatitis A virus cellular receptor 2 (HAVCR2), is a transmembrane glycoprotein of the TIM family of immune regulating molecules and plays an important role in the Th1-mediated immune response. TIM3 is expressed on the Th1 cells, CD8 T-cells, monocytes, and dendritic cells, but not on Th2 cells. TIM3 expressed by monocytes and dendritic cells facilitates phagocytosis of apoptotic cells and up-regulates cross-presentation of apoptotic cell-associated antigens through interaction with phosphatidylserine. Engagement of TIM3 by its ligand galectin-9 induces a range of immunosuppressive functions which enhance immune tolerance and inhibit anti-tumor immunity. Stimulation of TIM3 with an agonistic antibody promotes inflammation through the activation of innate immune cells. TIM3 is also regarded as a potential target molecule for immunotherapy. TIM3 and programmed cell death 1 (PD-1) as two important coinhibitory regulators of T cell responses, have been implicated with the T-cell dysfunction or exhaustion associated with chronic HBV infection including HBV-related HCC.</p> <p>Synonym: Hepatitis A virus cellular receptor 2 homolog, HAVcr-2, T-cell immunoglobulin and mucin domain-containing protein 3, T-cell immunoglobulin mucin receptor 3, T-cell membrane protein 3, Tim3, Timd3</p>
Molecular Weight:	46.3 kDa
UniProt:	<a href="#">Q8VIM0</a>
Pathways:	<a href="#">Regulation of Lipid Metabolism by PPARalpha</a> , <a href="#">Cancer Immune Checkpoints</a>


## Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

Handling

samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.