

Datasheet for ABIN7320866

Coxsackie Adenovirus Receptor Protein (His tag,AVI tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	Coxsackie Adenovirus Receptor (CXADR)
Origin:	Mouse
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Coxsackie Adenovirus Receptor protein is labelled with His tag,AVI tag.

Product Details

Purpose:	Recombinant Mouse CXADR/CAR Protein (AVI & His Tag)
Sequence:	Leu20-Gly237
Characteristics:	Biotinylated Recombinant Mouse CXADR is produced by our Mammalian expression system and the target gene encoding Leu20-Gly237 is expressed with a Avi, 6Histag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	Coxsackie Adenovirus Receptor (CXADR)
Alternative Name:	CXADR/CAR (CXADR Products)
Background:	Background: Coxsackievirus and adenovirus receptor homolog(CAR), also known as CXADR, is a type I transmembrane glycoprotein that belongs to the CTX family of the Ig superfamily. CXADR has monomer subunit that interacts with LNX, BAIAP1, DLG4, PRKCABP, TJP1 and

Target Details

CTNNB1. It also interacts with MPDZ and JAML. CXADR composed of of the epithelial apical junction complex that may function as a homophilic cell adhesion molecule and is essential for tight junction integrity. CXADR also involved in transepithelial migration of leukocytes through adhesive interactions with JAML a transmembrane protein of the plasma membrane of leukocytes. The interaction between both receptors also mediates the activation of gamma-delta T-cells, a subpopulation of T-cells residing in epithelia and involved in tissue homeostasis and repair. Upon epithelial CXADR-binding, JAML induces downstream cell signaling events in gamma-delta T-cells through PI3-kinase and MAP kinases. It results in proliferation and production of cytokines and growth factors by T-cells that in turn stimulate epithelial tissues repair.

Synonym: Coxsackievirus and adenovirus receptor homolog,CAR,Cxadr,CVB3 BP,MCVADR

Molecular Weight: 26.7 kDa

UniProt: [P97792](#)

Pathways: [Cell-Cell Junction Organization](#)

Application Details

Restrictions: For Research Use only

Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.