antibodies -online.com





SIGLEC10 Protein (Fc Tag)





Overview

Overview	
Quantity:	50 μg
Target:	SIGLEC10
Origin:	Mouse
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SIGLEC10 protein is labelled with Fc Tag.
Product Details	
_	

Purpose:	Recombinant Mouse Siglec-10 (C-Fc)
Sequence:	Met19-Lys543
Characteristics:	Recombinant Mouse Sialic Acid-binding Ig-like Lectin 10 is produced by our Mammalian expression system and the target gene encoding Met19-Lys543 is expressed with a Fc tag at the C-terminus.
Purity:	>95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	SIGLEC10
Alternative Name:	Siglec-10 (SIGLEC10 Products)
Background:	Background: Siglecs (sialic acid binding Ig-like lectins) are I-type lectins that belong to the
	immunoglobulin superfamily. They are characterized by an N-terminal Ig-like V-type domain

which mediates sialic acid binding, followed by a varying number of Ig-like C2-type domains. Siglecs 5-11 constitute the CD33/Siglec-3 related group, and are differentially expressed in the hematopoietic system. Siglec-G is the apparent ortholog of human Siglec-10. We describe here a novel member of the siglec protein family that shares a similar structure including five Ig-like domains, a transmembrane domain, and a cytoplasmic tail containing two ITIM-signaling motifs. Siglec-10 was identified through database mining of an asthmatic eosinophil EST library. Siglec-10 binds sialated proteins and lipids in alpha 2,3 or alpha 2,6 linkage and shows a preference for GT1b gangliosides. This binding can be modulated by cis interactions of Siglec-10 with sialated molecules expressed on the same cell. When tyrosine phosphorylated, the cytoplasmic ITIMs interact with phosphatases SHP-1 and SHP-2 to propagate inhibitory signals. The Siglec-10-VAP-1 interaction seems to mediate lymphocyte adhesion to endothelium and has the potential to modify the inflammatory microenvironment via the enzymatic end products.

Synonym: SIGLEC10, SiglecG, Siglec-G, MGC126774, PRO940, Siglec10, SLG2, sialic acid-binding Ig-like lectin 10, Siglec-10, siglec-like gene 2, Siglec-like protein 2, SLG2sialic acid binding Ig-like lectin 10 Ig-like lectin 7

Molecular Weight:

85.6 kDa

UniProt:

Q80ZE3

Application Details


Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.