antibodies .- online.com







CD70 Protein (His tag)



Image



Overview

Quantity:	50 µg
Target:	CD70
Origin:	Mouse
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD70 protein is labelled with His tag.
Product Details	

Purpose:	Recombinant Mouse CD27 Ligand/TNFSF7/CD70 (N-10His)
Sequence:	Gln47-Pro195
Characteristics:	Recombinant Mouse CD27 Ligand/CD70 is produced by our Mammalian expression system and the target gene encoding Gln47-Pro195 is expressed with a 10His tag at the N-terminus.
Purity:	>95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	CD70
Alternative Name:	CD27 Ligand/TNFSF7/CD70 (CD70 Products)
Background:	Background: CD27 Ligand (CD70) is a 30 kDa type 2 transmembrane glycoproteinis that belongs to the tumor necrosis factor (TNF) ligand family. Human CD27 Ligand extracellular
	region is 64 % aa identical to mouse and rat CD27 Ligand extracellular regions. This cytokine is

a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis.

Synonym: CD70 antigen, CD70, CD27 ligand, Tumor necrosis factor ligand superfamily member 7, CD27LG, TNFSF7, CD27L

Molecular Weight: 19 kDa
UniProt: Q05A52


Pathways: Cancer Immune Checkpoints

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.