

Datasheet for ABIN7320946
TNFRSF4 Protein (His tag)[Go to Product page](#)

1 Image

Overview

Quantity:	50 µg
Target:	TNFRSF4
Origin:	Cynomolgus
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TNFRSF4 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Cynomolgus OX40/TNFRSF4 Protein (His Tag)
Sequence:	Gly2-Pro512
Characteristics:	Recombinant Cynomolgus OX40L Receptor is produced by our Mammalian expression system and the target gene encoding Lys28-Ala214 is expressed with a 6His tag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	TNFRSF4
Alternative Name:	OX40/TNFRSF4 (TNFRSF4 Products)
Background:	Background: LYN contains one protein kinase domain, one SH2 domain, one SH3 domain, and the protein kinase domain plays an important role in its localization in the cell membrane. LYN is also thought to participate in signal transduction from cell surface receptors that lack

Target Details

intrinsic tyrosine kinase activity. In addition, LYN is associated with a number of cell surface receptors including the B cell antigen receptor and immunoglobulin E receptor (FcεRI).
Synonym: Tyrosine-protein kinase Lyn (LYN), Lck/Yes-related novel protein tyrosine kinase, V-src-1 Yamaguchi sarcoma viral related oncogene homolog, p53Lyn, p56Lyn and JTK8,

Molecular Weight:	21.3 kDa
UniProt:	P07948
Pathways:	Production of Molecular Mediator of Immune Response , Cancer Immune Checkpoints


Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.