

Datasheet for ABIN7320993
CD86 Protein (CD86) (His tag)[Go to Product page](#)

1 Image

Overview

| | |
|-------------------------------|---|
| Quantity: | 50 µg |
| Target: | CD86 |
| Origin: | Rabbit |
| Source: | Human Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CD86 protein is labelled with His tag. |

Product Details

| | |
|------------------|--|
| Purpose: | Recombinant Rabbit B7-2/CD86 Protein (His Tag) |
| Sequence: | Ala23-Arg246 |
| Characteristics: | Recombinant Rabbit CD86 is produced by our Mammalian expression system and the target gene encoding Ala23-Arg246 is expressed with a 6His tag at the C-terminus. |
| Purity: | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin Level: | < 1.0 EU per µg as determined by the LAL method. |

Target Details

| | |
|-------------------|--|
| Target: | CD86 |
| Alternative Name: | B7-2/CD86 (CD86 Products) |
| Background: | Background: T-lymphocyte activation antigen CD86 (B7-2) is a glycosylated protein in the B7 family. B7 family members are transmembrane cell surface molecules that play important roles in immune activation and the maintenance of immune tolerance. It is highly expressed on |

Target Details

activated antigen presenting cells. CD86 involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. It may play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. It is expressed by activated B-lymphocytes and monocytes and promoted by MARCH8 and results in endocytosis and lysosomal degradation.

Synonym: T-lymphocyte activation antigen CD86, Activation B7-2 antigen, CD86


| | |
|-------------------|--|
| Molecular Weight: | 26.3 kDa |
| UniProt: | P42071 |
| Pathways: | TCR Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Cellular Response to Molecule of Bacterial Origin , Positive Regulation of Immune Effector Process , Activated T Cell Proliferation |

Application Details

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling

| | |
|------------------|---|
| Format: | Lyophilized |
| Reconstitution: | Please refer to the printed manual for detailed information. |
| Buffer: | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. |
| Storage: | 4 °C,-20 °C,-80 °C |
| Storage Comment: | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |



Western Blotting

Image 1.