antibodies - online.com







CD47 Protein (CD47) (Fc Tag)





Overview

Quantity:	50 µg
Target:	CD47
Origin:	Rhesus Monkey
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD47 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Rhesus macaque CD47/IAP Protein (Fc Tag)
Sequence:	Gln19-Pro139
Characteristics:	Recombinant Rhesus macaque CD47 is produced by our Mammalian expression system and the target gene encoding Gln19-Pro139 is expressed with a Fc tag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	CD47
Alternative Name:	CD47/IAP (CD47 Products)
Background:	Background: CD47(Integrin-Associated Protein,IAP) is a 40 ? 60 kDa variably glycosylated atypical member of the immunoglobulin superfamily. The ubiquitously expressed CD47 binds
	to SIRP family members on macrophages, neutrophils, and T cells. CD47 is involved in the

Target Details

increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The protein is also a receptor for the C-terminal cell-binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This protein has broad tissue distribution, and is reduced in expression on Rh erythrocytes.

Synonym: Leukocyte Surface Antigen CD47, Antigenic Surface Determinant Protein OA3,

Molecular Weight:

40.8 kDa

UniProt:

F7A802


Application Details

Comment:	60-90 kDa	
Restrictions:	For Research Use only	

Integrin-Associated Protein, IAP, Protein MER6, CD47, MER6

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris, 100 mM Glycine, pH 7.5.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.