



[Go to Product page](#)

Datasheet for ABIN7321058

## Podoplanin Protein (PDPN) (Fc Tag,ECD)

### 1 Image

#### Overview

Quantity:	200 µg
Target:	Podoplanin (PDPN)
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Podoplanin protein is labelled with Fc Tag,ECD.

#### Product Details

Purpose:	Recombinant Rat Podoplanin/PDPN Protein (ECD, Fc Tag)
Sequence:	Met1-Leu135
Characteristics:	A DNA sequence encoding the rat Pdpn (NP_062231.1) (Met1-Leu135) was expressed with the Fc region of human IgG1 at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg protein as determined by the LAL method.

#### Target Details

Target:	Podoplanin (PDPN)
Alternative Name:	Podoplanin/PDPN ( <a href="#">PDPN Products</a> )
Background:	Background: Podoplanin, also known as PDPN, is a type-I integral membrane glycoprotein with diverse distribution in human tissues. The physiological function of this protein may be related to its mucin-type character. The homologous protein in other species has been described as a

## Target Details

---

differentiation antigen and influenza-virus receptor. The specific function of this protein has not been determined. Alternatively spliced transcript variants encoding different isoforms have been identified. PDPN is a mucin-type glycoprotein negatively charged by extensive O-glycosylation and a high content of sialic acid, which expresses the adhesive property. It is selectively expressed in lymphatic endothelium as well as lymphangiomas, Kaposi sarcomas, and in a subset of angiosarcomas with probable lymphatic differentiation. PDPN may contribute to form odontoblastic fiber or function as the anchorage to the tooth development and in proliferating epithelial cells of cervical loop and apical bud. The intensity of podoplanin expression is negatively correlated with the expression of CD34 and factor VIII. Podoplanin would be useful as a diagnostic marker for epithelioid hemangioendothelioma in liver tumors. Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy  
Synonym: Gp38;Ots8

---

Molecular Weight: 38.6 kDa

---

NCBI Accession: [NP\\_062231](#)

---

Pathways: [Dicarboxylic Acid Transport](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

---

Reconstitution: Please refer to the printed manual for detailed information.

---


Buffer: Lyophilized from sterile PBS, pH 7.4

---

Storage: 4 °C, -20 °C, -80 °C

---

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.